



Inland Counties Emergency Medical Agency

1425 South D Street, San Bernardino, CA 92415-0060 ■ (909) 388-5823 ■ Fax (909) 388-5825 ■ www.icema.net

Serving San Bernardino, Inyo, and Mono Counties
Tom Lynch, EMS Administrator
Reza Vaezazizi, MD, Medical Director

DATE: September 3, 2021

TO: EMS Providers - ALS, LALS, BLS, EMS Aircraft
Hospital CEOs, ED Directors, Nurse Managers and PLNs
EMS Training Institutions and Continuing Education Providers
Inyo, Mono and San Bernardino County EMCC Members
Medical Advisory Committee (MAC) Members
Systems Advisory Committee (SAC) Members

FROM: Tom Lynch
EMS Administrator

Reza Vaezazizi, MD
Medical Director

SUBJECT: IMPLEMENTATION OF POLICIES/PROTOCOLS EFFECTIVE OCTOBER 1, 2021

The revised policies/protocols listed below are effective October 1, 2021.

ICEMA Reference Number and Name

4060R2 Fireline EMT-P
4110 Trauma Critical Care System Designation (NEW)
6020R1 Responsibility for Patient Management
6100 EMS Fellow Field Response (NEW)
7010R3 Standard Drug and Equipment List - BLS/LALS/ALS
7020R2 Standard Drug and Equipment List - EMS Aircraft
8040 Transport of Patients (Inyo County Only) (DELETE)
8130R3 Assess and Refer Response Plan (San Bernardino County Only)
11010R3 Medication - Standard Orders
14110R1 Psychiatric/Behavioral Emergencies - Adult

Please insert and replace the enclosed policies/protocols and the Table of Contents in the Policy and Protocol Manual with the updated documents. The ICEMA policies and protocols can also be found on ICEMA's website at www.ICEMA.net under the Policy and Protocol Manual section.

If you have any questions, please contact Loreen Gutierrez, RN, Specialty Care Coordinator, at (909) 388-5803 or via e-mail at loreen.gutierrez@cao.sbcounty.gov.

TL/RV/jlm

Enclosures

c: File Copy

BOARD OF DIRECTORS

Col. Paul Cook (Ret.)
First District

Janice Rutherford
Second District

Dawn Rowe
Vice Chair
Third District

Curt Hagman
Chairman
Fourth District

Joe Baca, Jr.
Fifth District

Leonard X. Hernandez
Chief Executive Officer

POLICIES/PROTOCOLS CHANGES EFFECTIVE OCTOBER 1, 2021

Reference #	Name	Changes
DELETIONS		
8040	Transport of Patients (Inyo County Only)	Removed policy for Transport of Patients (Inyo County Only).
NEW		
4110	Trauma Critical Care System Designation	New policy for Trauma designation.
6100	EMS Fellow Field Response	New policy EMS Fellow Field Response.
CHANGES		
4060R2	Fireline EMT-P	Addition of Naloxone (Narcan).
6020R1	Responsibility for Patient Management	Clarified the transfer of care from an AEMT or EMT-P to an EMT.
7010R3	Standard Drug and Equipment List - BLS/LALS/ALS	Addition of Naloxone (Narcan) nasal spray. Moved CPAP from Optional to Non-Exchange Airway/Suction Equipment.
7020R2	Standard Drug and Equipment List - EMS Aircraft	Addition of Naloxone (Narcan) nasal spray.
8130R3	Assess and Refer Response Plan (San Bernardino County Only)	Updated policy name. Updated policy from an emergency surge response plan to a response plan that is active during normal operations.
11010R3	Medication - Standard Orders	Addition of Naloxone (Narcan) 4 mg IN loading dose for suspected Adult Fentanyl overdose.
14110R1	Psychiatric/Behavioral Emergencies - Adult	Changed the term excited delirium to potentially fatal and dangerous agitation.

TABLE OF CONTENTS

SERIES	OLD #	ADMINISTRATIVE POLICIES
1000		CREDENTIALING (EMT, AEMT, EMT-P, MICN)
1010	1100	AEMT Certification
1020	1030	EMT Certification
1030	1040	EMT-P Accreditation
1040R1	1050	MICN Authorization - Base Hospital, Administrative, Flight Nurse, Critical Care Transport
1050	1110	RCP Authorization
1060	1070	EMT/AEMT Incident Investigation, Determination of Action, Notification, and Administrative Hearing Process
1070	1090	Criminal History Background Checks (Live Scan)
1080	1120	EMT-P Student Field Internship Requirements
2000		EDUCATION
2010	3020	Continuing Education Provider Requirements
2020	3030	EMT Continuing Education Requirements
2030	3050	Public Safety First Aid Training Program Approval
2040	3060	Public Safety Optional Skills Course Approval
2050	3070	Tactical Casualty Care Training Programs and Courses
3000		GENERAL POLICIES
3010	5010	Licensure Changes - 911 Receiving Hospitals
3020R1	5020	Base Hospital Designation
3030	5030	Adoption of Policies and Protocols
3040R1	5040	Radio Communication
3050	7030	Controlled Substance
3060	5080	Ground Based Ambulance Rate Setting (San Bernardino County)
4000		SPECIALTY CARE PROGRAMS AND SPECIALTY SERVICE PROVIDER POLICIES
4010	6010	Paramedic Vaccination
4020	6170	ChemPack Deployment
4030	6060	Specialty and Optional Scope Program Approval
4040	6070	ST Elevation Myocardial Infarction Critical Care System Designation (San Bernardino County Only)
4050R1	6080	EMT-P Blood Draw for Chemical Testing at the Request of a Peace Officer
4060R2	6090	Fireline EMT-P
4070R1	6100	Stroke Critical Care System Designation (San Bernardino County Only)
4080R1	6110	Tactical Medicine for Special Operations
4090	6120	Emergency Medical Dispatch Center Requirements (San Bernardino County Only)
4100	6130	Medical Priority Dispatch Minimum Response Assignments for Emergency Medical Dispatch (EMD) Categories
4110	New	Trauma Critical Care System Designation

TABLE OF CONTENTS

SERIES	OLD #	EMS SYSTEM POLICIES
5000		DATA COLLECTION
5010	2020	ICEMA Abbreviation List
5020	2030	Minimum Documentation Requirements for Transfer of Patient Care
5030	2040	Requirements for Patient Care Reports
5040	2050	Requirements for Collection and Submission of EMS Data
6000		GENERAL OPERATIONAL POLICIES
6010	9020	Physician on Scene
6020R1	9030	Responsibility for Patient Management
6030	9040	Reporting Incidents of Suspected Abuse
6040	9050	Organ Donor Information
6050	9060	Local Medical Emergency
6060	9070	Patient Restraints
6070	9080	Care of Minors in the Field
6080	9090	Patient Refusal of Care - Adult
6090R1	9110	Treatment and Transportation Recommendations of Patients with Emerging Infectious Diseases
6100	New	EMS Fellow Field Response
7000		DRUG AND EQUIPMENT LISTS
7010R3	No Change	Standard Drug and Equipment List - BLS/LALS/ALS
7020R2	No Change	Standard Drug and Equipment List - EMS Aircraft
8000		RESPONSE, TRANSPORT, TRANSFER AND DIVERSION POLICIES
8010	No Change	Interfacility Transfer Guidelines
8020	No Change	Specialty Care Transport
8030	8050	Transport of Patients (BLS)
8040	8140	Transport of Patients (Inyo County Only) (DELETE)
8050	8060	Requests for Ambulance Redirection and Hospital Diversion (San Bernardino County Only)
8060	15050	Hospital Emergency Response Team (HERT)
8070R1	5070	Medical Response to Hazardous Materials/Terrorism Incident
8080R1	5050	Medical Response to a Multiple Casualty Incident
8090R1	5050I	Medical Response to a Multiple Casualty Incident (Inyo and Mono Counties)
8100	8150	Ambulance Patient Offload Delay (APOD)
8110	New	EMS Aircraft Utilization (San Bernardino County Only)
8120R1	New	Assess and Refer (San Bernardino County Only) (DELETE)
8130R3	New	Assess and Refer Response Plan (San Bernardino County Only)
9000		CONTINUATION OF CARE AND DESTINATION POLICIES
9010	8120	Continuation of Care (San Bernardino County Only)
9020R1	8090	Continuation of Trauma Care (Fort Irwin)
9030	8130	Destination
9040R1	15030	Trauma Triage Criteria

TABLE OF CONTENTS

10000		PILOT PROJECTS AND TRIAL STUDIES
10010	8160	Emergency Medical Transport of Police Dogs Pilot Project (San Bernardino County Only)
10020	6150	Trial Study Participation
11000		STANDARD ORDERS
11010R3	7040	Medication - Standard Orders
11020R2	10190	Procedure - Standard Orders

TABLE OF CONTENTS

SERIES	OLD #	PATIENT CARE PROTOCOLS
12000		GENERAL PATIENT CARE
12010R1	9010	Patient Care Guidelines
12020	15040	Glasgow Coma Scale
13000		ENVIRONMENTAL EMERGENCIES
13010	No Change	Poisonings
13020	No Change	Heat Related Emergencies
13030R1	No Change	Cold Related Emergencies
13040	No Change	Nerve Agent Antidote Kit (Training, Storage and Administration)
13050	11150	Smoke Inhalation/CO Exposure/Suspected Cyanide Toxicity
14000		TREATMENT PROTOCOLS
14010R2	11010	Respiratory Emergencies - Adult
14020R1	11020	Airway Obstruction - Adult
14030	11040	Bradycardias - Adult
14040	11050	Tachycardias - Adult
14050R1	11070	Cardiac Arrest - Adult
14060R1	11080	Altered Level of Consciousness/Seizures - Adult
14070R2	11100	Burns - Adult (15 years of age and older)
14080	11110	Stroke Treatment - Adult
14090R3	15010	Trauma - Adult (15 years of age and older)
14100	11140	Pain Management - Adult
14110R1	11130	Psychiatric/Behavioral Emergencies - Adult
14120	14010	Respiratory Emergencies - Pediatric (Less than 15 years of age)
14130	14020	Airway Obstruction - Pediatric (Less than 15 years of age)
14140	14030	Allergic Reactions - Pediatric (Less than 15 years of age)
14150	14040	Cardiac Arrest - Pediatric (Less than 15 years of age)
14160R1	14050	Altered Level of Consciousness - Pediatric (Less than 15 years of age)
14170	14060	Seizure - Pediatric (Less than 15 years of age)
14180R2	15020	Trauma - Pediatric (Less than 15 years of age)
14190	14070	Burns - Pediatric (Less than 15 years of age)
14200	14090	Newborn Care
14210	14080	Obstetrical Emergencies
14220	9120	Nausea and Vomiting
14230	9130	Shock (Non-Traumatic)
14240	11060	Suspected Acute Myocardial Infarction (AMI)
14250	12010	Determination of Death on Scene
14260	12020	End of Care and Decisions
14270	11120	Ventricular Assist Device (VAD)
15000		PUBLIC SAFETY FIRST AID
15010	16010	Allergic Reaction and Anaphylaxis (Authorized Public Safety Personnel)
15020	16020	Nerve Agent Exposure (Authorized Public Safety Personnel)
15030	16030	Opioid Overdose (Authorized Public Safety Personnel)
15040	16040	Respiratory Distress (Authorized Public Safety Personnel)
15050	16050	Optional Skills and Medications (Authorized Public Safety Personnel)
15060	16060	Public Safety AED Service Provider

TABLE OF CONTENTS

SERIES	OLD #	APPENDIX
16000		MISCELLANEOUS
16010	New	Definitions



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 4060R2
Effective Date: 10/01/21
Supersedes: 10/01/20
Page 1 of 3

FIRELINE EMT-P

I. PURPOSE

To provide guidance and medical oversight for an ICEMA paramedic (EMT-P) deployed to function as a fireline paramedic (FEMP).

This protocol is for use by authorized FEMPs during fire suppression activities and treatment of fire suppression personnel only.

II. REQUIREMENTS

- Must be a currently licensed EMT-P in California.
- Must be currently accredited EMT-P in the ICEMA region.
- Must be currently employed by an ICEMA approved ALS provider.
- The FEMP will follow FIRESCOPE FEMP ICS 223-11 Position Manual and all other ICS protocols.
- The FEMP will check in and obtain briefing from the Logistics Section Chief or the Medical Unit Leader, if established. Briefing will include current incident situation, anticipated medical needs, and local emergency medical system orientation.
- The FEMP will provide emergency medical treatment to personnel operating on the fireline.
- The FEMP will follow ICEMA prior to contact protocols if unable to contact the assigned base hospital.
- The FEMP may not perform skills outside of the ICEMA scope of practice.

III. PROCEDURE

- The EMS provider will notify ICEMA of the deployment of the FEMP to an incident. Use the Fireline Paramedic (FEMP) Deployment Notification form, which is on the ICEMA website at ICEMA.net.
- The FEMP will carry inventory in the advanced life support (ALS) pack as per the below inventory list (see Section IV. Fireline EMT-P (ALS) Pack Inventory). Inventory will be supplied and maintained by the employing ALS provider. Additional items for restock should also be maintained and secured in a vehicle or in the Medical Unit trailer.
- Incident Medical Units may not have the capability of resupplying controlled substances (narcotics). EMS providers should stock sufficient quantities of medical supplies and medications, especially controlled substance medications, to assure adequate supplies and medications.
- Narcotics must be under double lock and maintained on the FEMP person or secured in his/her vehicle at all times as per the ICEMA Drug and Equipment List.

- FEMP may carry an inventory of controlled substances (i.e., Fentanyl, Ketamine and Midazolam) if authorized by the employing ALS provider's Medical Director. The authorizing Medical Director is responsible to assure full compliance with all federal and state laws relating to purchase, storage and transportation of controlled substances. Only controlled substances approved for use in the ICEMA region may be carried and their use must be in accordance with current ICEMA patient care protocols.
- Radio communication failure protocols will not be used. Prior to base hospital contact protocols will be followed. If further treatment is needed, radio contact with the base hospital should be established as soon as possible.
- Documentation of patient care must follow ICEMA protocol utilizing the electronic patient care report (ePCR), if available, or a paper O1A form. All PCRs will be reviewed by the ALS provider and ICEMA for quality improvement (QI) purposes.
- A FEMP will be paired with a fireline EMT (FEMT) or another FEMP who will assist with basic life support (BLS) treatment and supplies.

IV. FIRELINE EMT-P (ALS) PACK INVENTORY

Minimum Requirements: The weight of the pack will dictate if the EMT-P chooses to carry additional ALS supplies.

MEDICATIONS/SOLUTIONS

Medications/Solutions	ALS
Albuterol Solution 2.5 mg Handheld Nebulizer or Multidose Inhaler	4
Atropine Sulfate 1 mg	2
Ipratropium Bromide Solution 0.5 mg Handheld Nebulizer or Multidose Inhaler	4
Lidocaine 100 mg IV pre-load	2
Aspirin 80 mg chewable	1 bottle
Dextrose 10%/250 ml (D10W 25 gm) IV/IO Bolus	1
Diphenhydramine 50 mg	4
Epinephrine 1: 10,000 1 mg	2
Epinephrine 1: 1000 1 mg	4
Glucagon 1 mg	1
Naloxone 2 mg	2
Nitroglycerin - Spray 0.4 metered dose and/or tablets (tablets to be discarded 90 days after opening)	1 (equivalent of 10 patient doses)
Saline 0.9% IV 1000 ml may be divided in two 500 ml bags or four 250 ml bags.	
Tranexamic Acid (TXA) 1 gm	1

CONTROLLED SUBSTANCE MEDICATIONS

Controlled Substance Medications MUST BE DOUBLED LOCKED	ALS
Midazolam	20 mg
Fentanyl (amount determined by the medical director)	200 - 400 mcg
Ketamine	120 - 1000 mg

ALS AIRWAY EQUIPMENT

Airway Equipment	ALS
Endotracheal Tubes - 6.0, 7.0 and/or 7.5 cuffed with stylet	1 each
Laryngeal blades - #0, #1, #2, #3, #4 curved and/or straight	1 each
Laryngoscope handle with batteries - or 2 disposable handles	1 each
ET Tube holder	1
End Tidal CO2 Detector	1
Needle Cricothyrotomy Kit	1
Needle Thoracostomy Kit	1

IV/MEDICATION ADMINISTRATION SUPPLIES

IV/Medication Administration Supplies	ALS
IV administration set macro drip	2
Venaguard	2
Alcohol preps	6
Betadine swabs	4
Tourniquet	2
Razor	1
Tape	1
IV catheters - 14, 16, 18 and 20 gauge	2
10 cc syringe	2
1 cc TB syringe	2
18 gauge needle	4
25 gauge needle	2

MISCELLANEOUS EQUIPMENT

Miscellaneous	ALS
Sharps container	1
Narcotic storage per protocol	
FEMP pack inventory sheet	1
Patient care report or ePCR (Toughbook)	
AMA forms	3

Equipment	ALS
Compact AED or compact monitor defibrillator combination	
Appropriate cardiac pads	
Pulse oximetry (optional)	
Glucometer, test strips and lancets	4

The BLS pack and supplies will be carried by the FEMT or accompanying FEMP. Personal items and supplies cannot be carried in either the ALS pack or the BLS pack.



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 4110 (NEW)

Effective Date: 10/01/21

Supersedes: NA

Page 1 of 4

TRAUMA CRITICAL CARE SYSTEM DESIGNATION

I. PURPOSE

To establish standards for the designation of an acute care hospital as a Trauma Receiving Center. These standards were developed to ensure patients who access the 9-1-1 system, and meet the defined Trauma triage criteria, are transported to a Trauma Receiving Center.

II. POLICY/PROCEDURE

Hospital requirements for Inland Counties Emergency Medical Agency (ICEMA) Trauma Receiving Center designation:

- Must be a full service general acute care hospital approved by ICEMA as a receiving hospital.
- Must have basic or comprehensive emergency services with special permits.
- Must be verified by the American College of Surgeons (ACS) as a Level I - III Trauma Receiving Center. Level IV Trauma Receiving Centers must remain in compliance with the current ACS standards.
- Must be in compliance with all requirements listed in California Code of Regulations, Title 22, Division 9, Chapter 7 - Trauma Critical Care System Regulations.

III. STAFFING REQUIREMENTS

The hospital will have the following positions filled prior to becoming a Trauma Receiving Center:

- Trauma Medical Directors

A qualified board-certified physician by the American Board of Medical Specialties (ABMS) as defined by the local EMS agency (LEMSA) and designated by the hospital that is responsible for the Trauma Receiving Center program, performance improvement, and patient safety programs related to a trauma critical care system.

- Emergency Department Trauma Representative

A qualified board certified emergency medicine physician with active privileges to practice in the emergency department that will participate in the Trauma Receiving Center program.

- Trauma Program Manager

The hospital shall designate a Trauma Program Manager who is responsible for monitoring and evaluating trauma patients. This includes participation in performance improvement and patient safety programs related to a trauma critical care system. The Trauma Program Manager must be trained or certified in critical care nursing and have continuing education in trauma physiology or at least has two (2) years dedicated trauma patient management experience.

- Trauma Team

A multidisciplinary team responsible for the initial resuscitation and management of the trauma patient.

- On-Call Physician Consultants and Staff

On-call physicians consultants and staff must be promptly available when notified. A daily roster must include the following on-call physician consultants and staff:

- Trauma Service: Must be promptly available, maximum trauma response time 15 minutes. Trauma surgeons must have privileges in general surgery and must be dedicated to a single Trauma Receiving Center while on duty (Level I and II).
- Neurosurgery Service: Must be promptly available for all traumatic brain injury (TBI) and spinal cord injury patients and must be present and respond within 30 minutes (Level I and II).
- Orthopedic Service: Must be promptly available for consultation within 30 minutes when requested by the trauma team leader (Level I and II).
- Anesthesiology Services: Must be available within 30 minutes for emergency operations.
- Radiology Services: Qualified radiologists must be available within 30 minutes in person or by tele radiology for the interpretation of radiographs.
- An operating room must be adequately staffed and available within 15 minutes (Level I and II).

- Registrar

A registrar dedicated to the registry must be available to process the data capturing the ICEMA data sets and in compliance with the ACS registrar standards listed in the "Resources for Optimal Care of the Injured Patient" current manual (Level I and II).

IV. INTERNAL HOSPITAL POLICIES

- The hospital must have capabilities to provide trauma patient care 24 hours per day, seven (7) days per week, 365 days per year.
- A single call alert/communication system for notification of incoming trauma patients, available 24 hours per day, seven (7) days per week (i.e., in-house paging system).
- The internal hospital policy/process/guidelines shall include:
 - A process for the treatment and triage of simultaneously arriving trauma patients.
 - A process for activation of trauma patients.
 - Prompt acceptance of trauma patients from referral hospitals per ICEMA Reference #9010 - Continuation of Care Policy.

- Acknowledgement that trauma patients may **only** be diverted during the times of Internal Disaster in accordance to ICEMA Reference #8050 - Requests for Ambulance Redirection and Hospital Diversion (San Bernardino County Only).
- A written notification describing the event must be submitted to ICEMA within 24 hours.
- A Level IV Trauma Receiving Center must have a written transfer agreement with a Level I or II Trauma Receiving Center, Level I or II Pediatric Trauma Receiving Center, or other specialty care centers, for immediate transfer of those patients for whom the most appropriate medical care requires additional resources.

V. DATA COLLECTION

All required data elements shall be collected and entered in an ICEMA approved Trauma registry on a quarterly basis and submitted to ICEMA for review. Trauma registry data must be collected in compliance with the National Trauma Data Standards and submitted to the National Trauma Data Bank (NTDB).

VI. CONTINUOUS QUALITY IMPROVEMENT (CQI) PROGRAM

- Trauma Receiving Centers shall develop an on-going CQI program which monitors all aspect of treatment and management of trauma patients and identify areas needing improvement. The program must, at a minimum, monitor the following parameters:
 - Mortality with opportunity for improvement.
 - Mortality without opportunity for improvement.
 - Unanticipated mortality with opportunity for improvement.
 - Rates of under-triage and over-triage.
- Active participation in quarterly regional Trauma Audit Committee and the regional Trauma peer review process. This will include a review of selected medical records as determined by CQI indicators and a presentation of details to peer review committee for adjudication.
- Provide continuing education (CE) opportunities twice per year for emergency medical services (EMS) field personnel in assessment and management of trauma patients.
- Programs in place to promote public education efforts specific to trauma care.

VII. PERFORMANCE STANDARD

Compliance with all California State Regulations and the ACS verification services performance standards.

VIII. DESIGNATION

- ICEMA designation as a Level I - IV Trauma Receiving Center will be based on an evaluation of need and volume in the community. Designation will not be determined by current compliance with Title 22 and compliance/verification of ACS alone; however, the Level I, II, and III Trauma Receiving Centers must be verified at least at an equivalent designation level that is being requested.

- The Trauma Receiving Center applicant shall be designated after satisfactory review of written documentation, a potential site survey by ICEMA, and completion of a board approved contractual agreement between the hospital and ICEMA.
- Documentation of current hospital accreditation by the ACS verification services as a Level I - III Trauma Receiving Center.
 - Level IV Trauma Receiving Centers must comply with all ACS Level IV standards.
- Initial designation as a Trauma Receiving Center shall be in accordance with terms outlined in the contract agreement.
- Failure to comply with the ICEMA policy, approved contract agreement, or the criteria and performance standards outlined in this policy, may result in probation, suspension fines or rescission of Trauma Receiving Center designation.

IX. REFERENCES

<u>Number</u>	<u>Name</u>
8050	Requests for Ambulance Redirection and Hospital Diversion (San Bernardino County Only)
9010	Continuation of Care (San Bernardino County Only)



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 6020R1
Effective Date: 10/01/21
Supersedes: 03/01/20
Page 1 of 1

RESPONSIBILITY FOR PATIENT MANAGEMENT

I. PURPOSE

To define the responsibility for patient care management in the prehospital setting. Within the ICEMA region, in the event both public and private emergency medical services (EMS) field personnel arrive on the scene with the same qualifications, patient care management responsibility will rest with the first to arrive.

II. PROCEDURE

- An advanced emergency medical technician (AEMT) or paramedic (EMT-P) may transfer patient management responsibility to an emergency medical technician (EMT) for transportation, only under the following conditions:
 - The patient is stable for transport and no ALS measures have been initiated.
 - When operating under ICEMA Reference #8030 - Transport of Patients (BLS).
 - When operating under ICEMA Reference #8080 - Medical Response to a Multiple Casualty Incident.
 - When operating under ICEMA Reference #6050 - Local Medical Emergency.
- The base hospital should be contacted if at any time transfer of patient management responsibility is in question or for any patient not meeting the above criteria.

III. REFERENCES

<u>Number</u>	<u>Name</u>
6050	Local Medical Emergency
8030	Transport of Patients (BLS)
8080	Medical Response to a Multiple Casualty Incident



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 6100 (NEW)

Effective Date: 10/01/21

Supersedes: NA

Page 1 of 2

EMS FELLOW FIELD RESPONSE

I. PURPOSE

To establish criteria for approved EMS Fellows and EMS Fellowship Leadership to serve as direct medical control when present in the field.

An EMS Fellow is a licensed physician who is participating in an accredited postgraduate EMS Fellowship training program following successful completion of a residency program in emergency medicine.

Once the EMS Fellow completes the required field training, this policy will allow an EMS Fellow to assist and/or direct paramedics (EMT-Ps) personnel in advanced life support (ALS) procedures according to ICEMA policies and protocols.

This policy applies specifically to physicians performing in the role as an EMS Fellow or Fellowship Leadership on scene, and does not pertain either to physicians who present as bystander citizens on scene or to physicians who are part of an established EMS response element (i.e., tactical physician, aeromedical flight team, search and rescue team).

II. POLICY/PROCEDURE

- ICEMA, the participating provider, along with EMS Fellow and Fellowship Leadership will determine field schedule.
- EMS field personnel will be notified of the EMS Fellow's field schedule prior to arrival.
- EMT field personnel shall obtain proper identification from the EMS Fellow and Fellowship Leadership.
- The EMS Fellow and Fellowship Leadership have the authority to provide on-scene medical direction.
- EMS field personnel may receive orders from the EMS Fellow and/or Fellowship Leadership within the Paramedic Scope of Practice and in compliance with ICEMA policy.
- The base hospital does not need to be contacted for orders.
- All EMS Fellow and Fellowship Leadership orders must be consistent with ICEMA policies and protocols.
- The EMS Fellow and Fellowship Leadership may perform medical care and procedures at the scene of an emergency.

III. PATIENT DESTINATION

- EMS field personnel are required to notify the receiving hospital that they are inbound.
- Patient will be transported to the most appropriate hospital in accordance with ICEMA Reference #9030 - Destination.

IV. LIABILITY

- Liability insurance is the responsibility of the EMS Fellowship program.

V. REFERENCES

<u>Number</u>	<u>Name</u>
9030	Destination



**INLAND COUNTIES
EMERGENCY MEDICAL AGENCY
POLICY AND PROTOCOL MANUAL**

Reference No. 7010R3

Effective Date: 10/01/21

Supersedes: 06/01/21

Page 1 of 6

STANDARD DRUG AND EQUIPMENT LIST - BLS/LALS/ALS

Each ambulance and first responder unit shall be equipped with the following functional equipment and supplies. **This list represents mandatory items with minimum quantities** excluding narcotics, which must be kept within the range indicated. All expiration dates must be current. All packaging of drugs or equipment must be intact. No open products or torn packaging may be used.

All ALS (transport and non-transport) and BLS transport vehicles shall be inspected annually.

MEDICATIONS/SOLUTIONS

Exchanged Medications/Solutions	BLS	LALS	ALS Non-Transport	ALS Transport
Adenosine (Adenocard) 6 mg			1	1
Adenosine (Adenocard) 12 mg			2	2
Albuterol Aerosolized Solution (Proventil) - unit dose 2.5 mg		4 doses	4 doses	4 doses
Albuterol MDI with spacer		1 SPECIALTY PROGRAMS ONLY	1 SPECIALTY PROGRAMS ONLY	1 SPECIALTY PROGRAMS ONLY
Aspirin, chewable - 81 mg tablet		2	1 bottle	1 bottle
Atropine 1 mg preload			2	2
Calcium Chloride 1 gm preload			1	1
Dextrose 10% in 250 ml Water (D10W)		2	2	2
Diphenhydramine (Benadryl) 50 mg			1	1
Epinephrine 1 mg/ml 1 mg		2	2	2
Epinephrine 0.1 mg/ml 1 mg preload			4	4
Glucagon 1 mg		1	1	1
Glucose paste	1 tube	1 tube	1 tube	1 tube
Ipratropium Bromide Inhalation Solution (Atrovent) unit dose 0.5 mg			4	4
Irrigating Saline and/or Sterile Water (1000 cc)	2	1	1	2
Lidocaine 100 mg			3	3
Lidocaine 2% Intravenous solution			1	1
Magnesium Sulfate 10 gm			1	1
Naloxone (Narcan) 2 mg preload	2	2	2	2
Nitroglycerine (NTG) - Spray 0.4 mg metered dose and/or tablets (tablets to be discarded 90 days after opening)		2	1	2
Nitroglycerine Paste 2% - 1 gm packets, or Nitroglycerine Paste 2% - 30 gm tube, or Nitroglycerine Paste 2% - 60 gm tube				2 1 1
Normal Saline for Injection (10 cc)		2	2	2
Normal Saline 100 cc			1	2
Normal Saline 250 cc			1	1
Normal Saline 500 ml and/or 1000 ml		2000 ml	3000 ml	6000 ml
Ondansetron (Zofran) 4 mg Oral Disintegrating Tablets (ODT)			4	4
Ondansetron (Zofran) 4 mg IM/IV			4	4
Sodium Bicarbonate 50 mEq preload			2	2
Tranexamic Acid (TXA) 1 gm			2	2

Non-Exchange Controlled Substance Medications MUST BE DOUBLE LOCKED	BLS	LALS	ALS Non- Transport	ALS Transport
Fentanyl			200-400 mcg	200-400 mcg
Midazolam			20-40 mg	20-40 mg
Ketamine			120-1000 mg	120-1000 mg

AIRWAY/SUCTION EQUIPMENT

Exchanged Airway/Suction Equipment	BLS	LALS	ALS Non- Transport	ALS Transport
CPAP circuits - all manufacture's available sizes			1 each	2 each
End-tidal CO2 device - Pediatric and Adult (may be integrated into bag)			1 each	1 each
Endotracheal Tubes cuffed - 6.0 and/or 6.5, 7.0 and/or 7.5 and 8.0 and/or 8.5 with stylet			2 each	2 each
ET Tube holders - adult		1 each	1 each	2 each
Mask - Adult & Pediatric non-rebreather oxygen mask	2 each	2 each	2 each	2 each
Mask - Infant Simple Mask	1	1	1	1
Nasal cannulas - pediatric and adult	2 each	2 each	2 each	2 each
Naso/Orogastric feeding tubes - 5fr or 6fr, and 8fr			1 each	1 each
Naso/Orogastric tubes - 10fr or 12fr, 14fr, 16fr or 18fr			1 each	1 each
Nasopharyngeal Airways - (infant, child, and adult)	1 each	1 each	1 each	1 each
Needle Cricothyrotomy Device - Pediatric and adult or Needles for procedure 10, 12, 14 and/or 16 gauge			1 each 2 each	1 each 2 each
One way flutter valve with adapter or equivalent			1	1
Oropharyngeal Airways - (infant, child, and adult)	1 each	1 each	1 each	1 each
Rigid tonsil tip suction	1		1	1
Small volume nebulizer with universal cuff adaptor		2	2	2
Suction Canister	1		1	1
Suction catheters - 6fr, 8fr or 10fr, 12fr or 14fr	1 each		1 each	1 each
Ventilation Bags - Infant 250 ml Pediatric 500 ml (or equivalent) Adult	1 1 1	1 1 1	1 1 1	1 1 1
Water soluble lubricating jelly		1	1	1

Non-Exchange Airway/Suction Equipment	BLS	LALS	ALS Non- Transport	ALS Transport
Ambulance oxygen source -10 L / min for 20 minutes	1			1
CPAP - (must be capable of titrating pressure between 2 and 15 cm H ₂ O)			1	1
Flashlight/penlight	1	1	1	1
Laryngoscope blades - #0, #1, #2, #3, #4 curved and/or straight			1 each	1 each
Laryngoscope handle with batteries - or 2 disposable handles			1	1
Magill Forceps - Pediatric and Adult			1 each	1 each
Manual powered suction device		1		
Portable oxygen with regulator - 10 L /min for 20 minutes	1	1	1	1
Portable suction device (battery operated)	1		1	1
Pulse Oximetry device	(SEE OPTIONAL EQUIPMENT SECTION, PG. 5)	1	1	1
Stethoscope	1	1	1	1
Wall mount suction device	1 (BLS TRANSPORT ONLY)			1

IV/NEEDLES/SYRINGES/MONITORING EQUIPMENT

Exchanged IV/Needles/Syringes/Monitor Equipment	BLS	LALS	ALS Non- Transport	ALS Transport
Conductive medium or Pacer/Defibrillation pads			2 each	2 each
Disposable Tourniquets		2	2	2
ECG electrodes			20	20
EZ-IO Driver			1 each	1 each
EZ-IO Needles: 25 mm 45 mm			2 each 1 each	2 each 1 each
Glucose monitoring device with compatible strips and OSHA approved single use lancets	1	1	1	1
3-way stopcock with extension tubing			2	2
IV Catheters - sizes 14, 16, 18, 20, 22, 24		2 each	2 each	2 each
Macro drip Administration Set		3	3	3
Microdrip Administration Set (60 drops / cc)		1	1	2
Mucosal Atomizer Device (MAD) for nasal administration of medication	2	2	2	4
Pressure Infusion Bag (disposable)		1	1	1
Razors		1	2	2
Safety Needles - 20 or 21gauge and 23 or 25 gauge	2 each	2 each	2 each	2 each
Saline Lock Large Bore Tubing Needleless		2	2	2
Sterile IV dressing		2	2	2
Syringes w/wo safety needles - 1 cc, 3 cc, 10 cc catheter tip		2 each		

Exchanged IV/Needles/Syringes/Monitor Equipment	BLS	LALS	ALS Non-Transport	ALS Transport
Syringes w/wo safety needles - 1 cc, 3 cc, 10 cc, 20 cc, 60 cc catheter tip			2 each	2 each

Non-Exchange IV/Needles/Syringes/ Monitor Equipment	BLS	LALS	ALS Non-Transport	ALS Transport
12-lead ECG Monitor and Defibrillator with TCP and printout			1	1
Blood pressure cuff - large adult or thigh cuff, adult, child and infant (one of each size)	1	1	1	1
Capnography monitor and supplies, may be integrated in the cardiac monitor			1	1
Needle disposal system (OSHA approved)	1	1	1	1
Thermometer - Mercury Free with covers	1	1	1	1

OPTIONAL EQUIPMENT/MEDICATIONS

Non-Exchange Optional Equipment/ Medications	BLS	LALS	ALS Non-Transport	ALS Transport
AED/defib pads - Adult (1), Pediatric (1)	1 each	1 each		
Automatic CPR device (FDA approved)	1	1	1	1
Automatic transport ventilator (Specialty Program Only - ICEMA approved device)			1	1
Backboard padding	1	1	1	1
Buretrol			1	1
Chemistry profile tubes			3	3
Nerve Agent Antidote Kit (NAAK) - DuoDote or Mark I	3	3	3	3
EMS Tourniquet	1		1	1
Gum Elastic intubation stylet			2	2
Hemostatic Dressings *	1	1	1	1
IO Needles - Manual, Adult and Pediatric, Optional		Pediatric sizes only or EZ-IO needles and drivers	1 each	1 each
IV infusion pump			1	1
IV warming device		1	1	1
Manual IV Flow Rate Control Device			1	1
Manual powered suction device	1	1	1	1
Multi-lumen peripheral catheter			2	2
Needle Thoracostomy Kit (prepackaged)			2	2
Naloxone (Narcan) Nasal Spray 4 mg	2	2	2	2
Pulse Oximetry device	1			
Translaryngeal Jet Ventilation Device			1	1
Vacutainer			1	1

* Hemostatic Dressings

- Quick Clot, Z-Medica
 - Quick Clot, Combat Gauze LE
 - Quick Clot, EMS Rolled Gauze, 4x4 Dressing, TraumaPad
- Celox
 - Celox Gauze, Z-Fold Hemostatic Gauze

Celox Rapid, Hemostatic Z-Fold Gauze

- HemCon ChitoFlex Pro Dressing

NOTE:

- The above products are “packaged” in various forms (i.e., Z-fold, rolled gauze, trauma pads, 4”x4”pads) and are authorized provided they are comprised of the approved product.
- Hemostatic Celox Granules, or granules delivered in an applicator, are not authorized.

DRESSING MATERIALS/OTHER EQUIPMENT/SUPPLIES

Exchanged Dressing Materials/Other Equipment/Supplies	BLS	LALS	ALS Non-Transport	ALS Transport
Adhesive tape - 1 inch	2	2	2	2
Air occlusive dressing	1	1	1	1
Ankle and wrist restraints, soft ties acceptable	1		1	1
Antiseptic swabs/wipes	10	10	10	10
Bedpan or fracture pan	1 (BLS TRANSPORT UNITS ONLY)			1
Urinal	1 (BLS TRANSPORT UNITS ONLY)			1
Cervical Collars - Rigid Pediatric and Adult all sizes or	2 each	2 each	2 each	2 each
Cervical Collars - Adjustable Adult and Pediatric	2 each	2 each	2 each	2 each
Cold Packs	2	2	2	2
Emesis basin or disposable bags and covered waste container	1	1	1	1
Head immobilization device	2	2	2	2
OB Kit	1	1	1	1
Pneumatic or rigid splints capable of splinting all extremities	4	2	2	4
Providine/Iodine swabs/wipes or antiseptic equivalent		4	10	10
Roller bandages - 4 inch	6	3	3	6
Sterile bandage compress or equivalent	6	2	2	6
Sterile gauze pads - 4x4 inch	4	4	4	4
Sterile sheet for Burns	2	2	2	2
Universal dressing 10x30 inches	2	2	2	2

Non-Exchange Dressing Materials/Other Equipment/Supplies	BLS	LALS	ALS Non-Transport	ALS Transport
800 MHz Radio		1	1	1
Ambulance gurney	1 (BLS TRANSPORT UNITS ONLY)			1
Bandage shears	1	1	1	1
Blood Borne Pathogen Protective Equipment - (nonporous gloves, goggles face masks and gowns meeting OSHA Standards)	2	1	2	2

Non-Exchange Dressing Materials/Other Equipment/Supplies	BLS	LALS	ALS Non-Transport	ALS Transport
Pediatric Emergency Measuring Tape (Broselow, etc.)		1	1	1
Drinkable water in secured plastic container or equivalent	1 gallon			1 gallon
Long board with restraint straps	1	1	1	1
Pediatric immobilization board	1	1	1	1
Pillow, pillow case, sheets and blanket	1 set (BLS TRANSPORT UNITS ONLY)			1 set
Short extrication device	1	1	1	1
Straps to secure patient to gurney	1 set (BLS TRANSPORT UNITS ONLY)			1 set
Traction splint	1	1	1	1
Triage Tags - ICEMA approved	20	20	20	20



**INLAND COUNTIES
EMERGENCY MEDICAL AGENCY
POLICY AND PROTOCOL MANUAL**

Reference No. 7020R2
Effective Date: 10/01/21
Supersedes: 10/01/20
Page 1 of 3

STANDARD DRUG AND EQUIPMENT LIST - EMS AIRCRAFT

Each Aircraft shall be equipped with the following functional equipment and supplies. This list represents mandatory items with minimum quantities, to exclude narcotics, which must be kept within the range indicated. All expiration dates must be current. All packaging of drugs or equipment must be intact. No open products or torn packaging may be used.

MEDICATIONS/SOLUTIONS	AMOUNT
Adenosine (Adenocard) 6 mg	1
Adenosine (Adenocard) 12 mg	2
Albuterol Aerosolized Solution (Proventil) - unit dose 2.5 mg	3 doses
Aspirin, chewable - 81 mg tablet	1 bottle
Atropine 1 mg preload	2
Calcium Chloride 1 gm preload	1
Dextrose 10% in 250 ml Water (D10W) *	2
Diphenhydramine (Benadryl) 50 mg	1
Epinephrine 1 mg/ml 1 mg	2
Epinephrine 0.1 mg/ml 1mg preload	3
Glucagon 1 mg	1
Glucopaste	1 tube
Ipratropium Bromide Inhalation Solution (Atrovent) unit dose 0.5 mg	3
Lidocaine 100 mg	3
Lidocaine 2% Intravenous solution	1
Magnesium Sulfate 10 gms	1
Naloxone (Narcan) 2 mg preload	2
Nitroglycerin (NTG) - Spray 0.4 mg metered dose and/or tablets (tablets to be discarded 90 days after opening.)	1
Normal Saline for Injection (10 cc)	2
Normal Saline 250 ml	1
Normal Saline 500 ml and/or 1000 ml	2000 ml
Ondansetron (Zofran) 4 mg Oral Disintegrating Tablets (ODT)	4
Ondansetron (Zofran) 4 mg IM/ IV	4
Sodium Bicarbonate 50 mEq preload	2
Tranexamic Acid (TXA) 1 gm	2

CONTROLLED SUBSTANCE MEDICATIONS-MUST BE DOUBLE LOCKED	AMOUNT
Fentanyl	200-400 mcg
Midazolam	20-40 mg
Ketamine	120-1000 mg

AIRWAY/SUCTION EQUIPMENT	AMOUNT
Aircraft Oxygen source -10 L /min for 20 minutes	1
C-PAP circuits - all manufacture's available sizes	1 each
End-tidal CO2 device - pediatric and adult (may be integrated into bag)	1 each
Endotracheal Tubes cuffed - 6.0 and/or 6.5, 7.0 and/or 7.5 and 8.0 and/or 8.5 with stylet	2 each
ET Tube holders - adult	1 each
Flashlight/penlight	1
Laryngoscope handle with batteries - or 2 disposable handles	1
Laryngoscope blades - #0, #1, #2, #3, #4 curved and/or straight	1 each
Magill Forceps - Pediatric and Adult	1 each
Nasal Cannulas - infant, pediatric and adult	2 each
Naso/Orogastric tubes - 10fr or 12fr, 14fr, 16fr or 18fr	1 each

AIRWAY/SUCTION EQUIPMENT	AMOUNT
Naso/Orogastric feeding tubes - 5fr or 6fr, and 8fr	1 each
Nasopharyngeal Airways - infant, child, and adult	1 each
Needle Cricothyrotomy Device (Approved) - Pediatric and adult <i>or</i>	1 each
Needles for procedure 10, 12, 14 and/or 16 gauge	2 each
Non Re-Breather O ₂ Mask - Pediatric and Adult, Infant Simple Mask	2 each
One way flutter valve with adapter or equivalent	1
Oropharyngeal Airways - infant, child, and adult	1 each
Portable Oxygen with regulator - 10 L /min for 20 minutes	1
Portable suction device (battery operated) <i>and/or</i> Wall mount suction device	1 each
Pulse Oximetry device	1
Small volume nebulizer with universal cuff adaptor	1
Stethoscope	1
Suction catheters - 6fr, 8fr or 10fr, 12fr or 14fr	1 each
Ventilation Bags - Infant 250 ml, Pediatric 500 ml and Adult 1 L	1 each
Water soluble lubricating jelly	1
Ridged tonsil tip suction	1

IV/NEEDLES/SYRINGES/MONITORING EQUIPMENT	AMOUNT
12-Lead ECG Monitor and Defibrillator with TCP and printout	1
800 MHz Radio	1
Blood pressure cuff - large adult or thigh cuff, adult, child and infant	1 set
Capnography monitor and supplies, may be integrated in the cardiac monitor	1
Conductive medium <i>or</i> Adult and Pediatric Pacer/Defibrillation pads	2 each
ECG - Pediatric and Adult	20 patches
EZ IO Needles and Driver 25 mm <i>and</i>	2 each
45 mm	1 each
3-way stopcock with extension tubing	2
IO Needles - Manual, Adult and Pediatric, <u>Optional</u>	1 each
IV Catheters - sizes 14, 16, 18, 20, 22, 24	2 each
Glucose monitoring device	1
Macro drip Administration Set	3
Micro drip Administration Set (60 drops/ml)	1
Mucosal Atomizer Device (MAD) for nasal administration of medication	4
Needle disposal system (OSHA approved)	1
Pressure infusion bag	1
Safety Needles - 20 or 21 gauge and 23 or 25 gauge	2 each
Saline Lock	2
Syringes w/wo safety needles - 1 ml, 3 ml, 10 ml, 20 ml	2 each
Syringe - 60 ml catheter tip	2
Thermometer - Mercury free with covers	1

DRESSING MATERIALS/OTHER EQUIPMENT SUPPLIES	AMOUNT
Adhesive tape - 1 inch	2
Air occlusive dressing	1
Aircraft stretcher or litter system with approved FAA straps that allows for Axial Spinal Immobilization	1
Ankle and wrist restraints, soft ties acceptable	1
Antiseptic swabs/wipes	
Bandage shears	1
Blanket or sheet	2

DRESSING MATERIALS/OTHER EQUIPMENT SUPPLIES	AMOUNT
Blood Borne Pathogen Protective Equipment - (nonporous gloves, goggles face masks and gowns meeting OSHA Standards)	2
Cervical Collars - Rigid Pediatric & Adult all sizes <i>or</i> Cervical Collars - Adjustable Adult and Pediatric	1 each 1 each
Emesis basin or disposable bags and covered waste container	1
Head immobilization device	1
OB Kit	1
Pediatric Emergency Measuring Tape (Broselow, etc.)	1
Pneumatic or rigid splints capable of splinting all extremities	4
Providine/Iodine swabs/wipes or antiseptic equivalent	
Roller bandages - 4 inch	3
Sterile bandage compress or equivalent	6
Sterile gauze pads - 4x4 inch	4
Sterile Sheet for Burns	2
Traction splint	1
Universal Dressing 10x30 inches	2

OPTIONAL EQUIPMENT/MEDICATIONS	Amount
Automatic ventilator (Approved)	1
Backboard padding	1
BLS AED/defib pads	1
Chemistry profile tubes	3
Nerve Agent Antidote Kit (NAAK) - DuoDote or Mark I	3
D5W in bag	1
Hemostatic Dressing *	1
IV infusion pump	1
IV warming device	1
Manual powered suction device	1
Medical Tourniquet	1
Naloxone (Narcan) Nasal Spray 4 mg	2
Needle Thoracostomy Kit (prepackaged)	2
Pediatric immobilization board	1
Translaryngeal Jet Ventilation Device	1
Vacutainer	1

* Hemostatic Dressings

- Quick Clot, Z-Medica
 - Quick Clot, Combat Gauze LE
 - Quick Clot, EMS Rolled Gauze, 4x4 Dressing, TraumaPad
- Celox
 - Celox Gauze, Z-Fold Hemostatic Gauze
 - Celox Rapid, Hemostatic Z-Fold Gauze

NOTE:

- The above products are “packaged” in various forms (i.e., Z-fold, rolled gauze, trauma pads, and 4”x4” pads) and are authorized provided they are comprised of the approved product.
- Hemostatic Celox Granules, or granules delivered in an applicator, are not authorized.



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 8040
Effective Date: 03/01/20
Supersedes: 12/15/15
Page 1 of 2

TRANSPORT OF PATIENTS (INYO COUNTY ONLY)

I. PURPOSE

To provide guidelines for EMS field personnel for the transportation of patients in Inyo County.

II. POLICY

A. Ground Transport of Patients in Inyo County

- All patients originating in Independence (EOA 3) shall be transported to Northern Inyo Hospital (NIH) per ICEMA Reference #9030 - Destination.
- All patients originating in Olancho/Cartago (EOA 5 and 6) shall be transported to Ridgecrest Regional Hospital (RRH) per ICEMA Reference #9030 - Destination.
- Advanced Life Support (ALS) intercept may be used when available and only when the patient's condition requires a higher level of care.
- The receiving hospital shall be contacted as soon as possible according to ICEMA Reference #3040 - Radio Communication.
- Base hospital physician may override prior destination decision by paramedic (EMT-P) per ICEMA Reference #9030 - Destination.

NOTE: As a reference, Cottonwood Creek Bridge (halfway between Olancho/Cartago and Lone Pine) is the mid-point between NIH and RRH.

B. Special Considerations

All patients originating in Lone Pine (EOA 4) that require a higher level of care:

- An ALS flight crew (Sierra Lifeflight) may be requested if ALS care is required.
- Simultaneously base hospital contact shall be made to base hospital who will determine (in collaboration with the ground and flight crew) whether:
 - Patient is transported via ground to NIH with ALS flight crew.
 - Patient is transported via air to Bishop and then by ground to NIH.
 - Patient is transported by air or ground to a hospital outside the county.

C. Base Hospital Contact

- Base hospital contact is required according to ICEMA Reference #3040 - Radio Communication.
- All patients being considered for transport to hospitals other than NIH or RRH require NIH base hospital contact for medical control and destination decision.


- Patients requiring higher level of care such as that required by patient condition (trauma, stroke or STEMI), may be directed to a more distant facility by the base hospital.
- All patient destinations other than by ground transport require base hospital direction prior to transport.

D. Patient Documentation and Quality Improvement (QI/QA)

- EMS field personnel must complete an ICEMA approved electronic patient care report (ePCR) for all patients.
- All ePCRs will be reviewed as part of the EMS provider and base hospital review process.

III. REFERENCES

<u>Number</u>	<u>Name</u>
3040	Radio Communication
9030	Destination





INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 8130R3
Effective Date: 10/01/21
Supersedes: 12/10/20
Page 1 of 3

ASSESS AND REFER RESPONSE PLAN (San Bernardino County Only)

I. PURPOSE

To establish standards for the identification of patients whose condition does not require transport by 9-1-1 emergency ambulance. All 9-1-1 calls for EMS will receive an appropriate response, timely assessment, and appropriate patient care. If it is determined that the patient is stable, and does not require emergency department services EMS field personnel will assess patient and provide an appropriate alternative recommendation.

II. POLICY

- If the patient's condition is stable and meets assess and refer criteria EMS field personnel will provide the patient the following recommendation:
 - "It appears that you do not require immediate care in the emergency department. You should seek care with your regular healthcare provider, urgent care or clinic. If symptoms worsen seek medical help or re-contact 9-1-1."

III. GENERAL CONSIDERATIONS

- Transport all patients requiring immediate medical attention to the closest most appropriate hospital.
- EMS should not require patients that are being released from the scene to sign AMA on the Patient Care Record.
- Provide instructions that if symptoms worsen, patient should go to the emergency department, contact their healthcare provider, or re-contact 9-1-1.
- If the patient or guardian refuses the referral, the patient will be transported to the closest most appropriate hospital.

IV. PARAMEDIC ASSESS AND REFER DECISION MAKING PRINCIPLES

- Does the patient, parent, or guardian have Decision Making Capacity?
- Is EMS field personnel concerned with the patient's current medical condition?
- How likely is the patient to successfully navigate the provided referral?

V. ASSESS AND REFER CRITERIA

- The patient must meet all of the following criteria:
 - Parent or guardian is on scene if the patient is under 18 years of age (unless legally emancipated).
 - Has a Glasgow Coma Scale (GSC) of 15 or GCS is at patient's baseline.
 - Exhibits no clinical evidence of:
 - Altered level of consciousness

- Alcohol or drug ingestion that impairs decision making capacity
 - Abnormal or labored breathing or shortness of breath
 - Chest pain/discomfort of any kind
 - Hypoxia as indicated by low oxygen saturation
 - Significant tachycardia
 - Serious hemorrhage
- Exhibits evidence of Decision-Making Capacity sufficient to understand the nature of the medical condition as well as the risks and potential consequences of not seeking additional medical care from the provided recommendation.
 - The patient would benefit from the provided recommendation.
 - The patient is likely to successfully navigate the provided recommendation.
- If there is clinical evidence of a viral illness, the patient must meet all the following criteria:
 - Be stable.
 - Not under two (2) years of age, or over 65 years of age.
 - Does not have an underlying medical history.
 - For the COVID positive patient or PUI, assess for a referral to stay home, self-isolate, and seek follow-up treatment with a physician.

VI. DOCUMENTATION REQUIREMENTS

- Physical exam.
- Treatment provided.
- Patient, parent, or guardian is alert, oriented, and acting appropriately for their age.
- Indications that there were no signs of significant impairment due to drugs, alcohol, organic causes, or mental illness.
- Any other observations that indicate that the patient, parent, or guardian has impaired Decision-Making Capacity.
- Recommendation/referrals shall be documented utilizing the following four (4) step process:
 - That a recommendation/referral was offered.
 - What the recommendation/referral was that EMS field personnel provided.
 - The patient's understanding of the recommendation/referral.
 - The patient's plan based on the recommendation/referral of the EMS field personnel.
- The person(s), if any, who remained to look after the patient (the patient's "support system").

- The name of the interpreter utilized, if applicable.
- EMS field personnel will leave a referral card containing relevant community referral information with the patient.



**INLAND COUNTIES
EMERGENCY MEDICAL AGENCY
POLICY AND PROTOCOL MANUAL**

Reference No. 11010R3

Effective Date: 10/01/21

Supersedes: 06/15/21

Page 1 of 13

MEDICATION - STANDARD ORDERS

Medications listed in this protocol may be used only for the purposes referenced by the associated ICEMA Treatment Protocol.

For Nerve Agent Antidote Kit (NAAK) or medications deployed with the ChemPack see Appendix I (Page 12).

Adenosine (Adenocard) - Adult (ALS)

Stable narrow-complex SVT or Wide complex tachycardia:

Adenosine, 6 mg rapid IVP followed immediately by 20 cc NS bolus, and Adenosine, 12 mg rapid IVP followed immediately by 20 cc NS bolus if patient does not convert. May repeat one (1) time.

Reference #s 7010, 7020, 14040

Albuterol (Proventil) Aerosolized Solution - Adult (LALS, ALS)

Albuterol, 2.5 mg nebulized, may repeat two (2) times.

Reference #s 4060, 7010, 7020, 14010, 14070

Albuterol (Proventil) Metered-Dose Inhaler (MDI) - Adult (LALS, ALS - Specialty Programs Only)

Albuterol MDI, four (4) puffs every 10 minutes for continued shortness of breath and wheezing.

Reference #s 4060, 4080, 7010, 7020, 14120, 14140, 14190

Albuterol (Proventil) - Pediatric (LALS, ALS)

Albuterol, 2.5 mg nebulized, may repeat two (2) times.

Reference #s 7010, 7020, 14120, 14140, 14190

Albuterol (Proventil) Metered-Dose Inhaler (MDI) - Pediatric (LALS, ALS - Specialty Programs Only)

Albuterol MDI, four (4) puffs every 10 minutes for continued shortness of breath and wheezing.

Reference #s 4060, 4080, 7010, 7020, 14120, 14140, 14190

Aspirin, chewable (LALS, ALS)

Aspirin, 325 mg PO chewed (one (1) adult non-enteric coated aspirin) or four (4) chewable 81 mg aspirin.

Reference #s 4060, 4080, 5010, 7010, 7020, 14240

Atropine (ALS) - Adult

Atropine, 0.5 mg IV/IO. May repeat every five (5) minutes up to a maximum of 3 mg or 0.04 mg/kg.

Organophosphate poisoning:

Atropine, 2 mg IV/IO, repeat at 2 mg increments every five (5) minutes if patient remains symptomatic.

Reference #s 4060, 4080, 7010, 7020, 13010, 14030, 14260

Atropine - Pediatric (ALS)*Organophosphate poisoning - Pediatrics less than 14 years of age:*

Atropine, 0.05 mg/kg IV/IO not to exceed adult dose of 2 mg, repeat at 0.1 mg/kg increments every five (5) minutes if patient remains symptomatic.

Reference #s 4060, 4080, 7010, 7020, 13010

Calcium Chloride - Adult (ALS)*Calcium Channel Blocker Poisonings (base hospital order only):*

Calcium Chloride, 1 gm (10 ml of a 10% solution) IV/IO.

Reference #s 5010, 7010, 7020, 13010

For cardiac arrest with suspected hypocalcemia, hyperkalemia, hypermagnesemia or calcium channel blocker poisoning (base hospital order only):

Calcium Chloride, 1 gm (10 ml of a 10% solution) IV/IO.

Reference #s 7010, 7020, 14050

Calcium Chloride - Pediatric (ALS)*Calcium Channel Blocker Poisonings (base hospital order only):*

Calcium Chloride, 20 mg/kg IV/IO over five (5) minutes.

Reference #s 7010, 7020, 13010

Dextrose - Adult (LALS, ALS)*Hypoglycemia - Adult with blood glucose less than 80 mg/dL:*

Dextrose 10% /250 ml (D10W 25 gm) IV/IO Bolus

Reference #s 4060, 4080, 5010, 7010, 7020, 8010, 13020, 13030, 14040, 14060

Dextrose - Pediatric (LALS, ALS)*Hypoglycemia - Neonates (0 - 4 weeks) with blood glucose less than 35 mg/dL or pediatric patients (more than 4 weeks) with glucose less than 60 mg/dL:*

Dextrose 10%/250 ml (D10W 25 gm) 0.5 gm/kg (5 ml/kg) IV/IO

Reference #s 5010, 7010, 7020, 13020, 13030, 14150, 14160, 14170

Diphenhydramine - Adult (ALS)

Diphenhydramine, 25 mg IV/IO

Diphenhydramine, 50 mg IM

Reference #s 4060, 4080, 7010, 7020, 13010, 14010

Diphenhydramine - Pediatric (ALS)

Allergic reaction:

2 years to 14 years Diphenhydramine, 1 mg/kg slow IV/IO, not to exceed adult dose of 25 mg, **or**

Diphenhydramine, 2 mg/kg IM not to exceed adult dose of 50 mg IM.

Reference #s 7010, 7020, 14140

Epinephrine (1 mg/ml) - Adult (LALS, ALS)

Severe Bronchospasm, Asthma Attack, Pending Respiratory Failure, Severe Allergic Reactions:

Epinephrine, 0.3 mg IM. May repeat after 15 minutes one (1) time if symptoms do not improve.

Reference # 14010

Epinephrine (0.1 mg/ml) - Adult (ALS)

For persistent severe anaphylactic reaction:

Epinephrine (0.1 mg/ml), 0.1 mg slow IVP/IO. May repeat every five (5) minutes as needed to total dosage of 0.5 mg.

Reference # 14010

Cardiac Arrest, Asystole, PEA:

Epinephrine (0.1 mg/ml), 1 mg IV/IO.

Reference #s 4060, 4080, 5010, 7010, 7020, 14010, 14050, 14260

Epinephrine (0.01 mg/ml) - Adult (ALS)

Post resuscitation, persistent profound nontraumatic shock and hypotension (Push Dose Epinephrine):

Prepare Epinephrine 0.01 mg/ml solution by mixing 9 ml of normal saline with 1 ml of Epinephrine 0.1 mg/ml in a 10 ml syringe. Administer 1 ml every one (1) to five (5) minutes titrated to maintain SBP more than 90 mm Hg.

Reference #s 4060, 4080, 5010, 7010, 7020, 11010, 14050, 14230

Epinephrine (1 mg/ml) - Pediatric (LALS, ALS)

Severe Bronchospasm, Asthma Attack, Pending Respiratory Failure, Severe Allergic Reactions:

Epinephrine, 0.01 mg/kg IM not to exceed adult dosage of 0.3 mg.

Reference #s 4060, 5010, 7010, 7020, 14120, 14140

Epinephrine (0.1 mg/ml) - Pediatric (ALS)

Anaphylactic reaction (no palpable radial pulse and depressed level of consciousness):

Epinephrine (0.1 mg/ml), 0.01 mg/kg IV/IO, no more than 0.1 mg per dose. May repeat to a maximum of 0.5 mg.

Cardiac Arrest:

1 day to 8 years Epinephrine (0.1 mg/ml), 0.01 mg/kg IV/IO (do not exceed adult dosage)

9 to 14 years Epinephrine (0.1 mg/ml), 1.0 mg IV/IO

Newborn Care:

Epinephrine (0.1 mg/ml), 0.01 mg/kg IV/IO if heart rate is less than 60 after one (1) minute after evaluating airway for hypoxia and assessing body temperature for hypothermia.

Epinephrine (0.1 mg/ml), 0.005 mg/kg IV/IO every 10 minutes for persistent hypotension as a base hospital order or in radio communication failure.

Reference # 14200

Epinephrine (0.01 mg/ml) - Pediatric (ALS)

Post resuscitation, profound shock and hypotension (Push Dose Epinephrine):

Prepare Epinephrine 0.01 mg/ml solution by mixing 9 ml of normal saline with 1 ml of Epinephrine 0.1 mg/ml in a 10 ml syringe. Administer 0.1 ml/kg (do not exceed adult dosage), every one (1) to five (5) minutes. Titrate to maintain a SBP more than 70 mm Hg.

Reference #s 5010, 7010, 7020, 11010, 14150, 14230

Fentanyl - Adult (ALS)

Chest Pain (Presumed Ischemic Origin):

Fentanyl, 50 mcg slow IV/IO over one (1) minute. May repeat every five (5) minutes titrated to pain, not to exceed 200 mcg.

Fentanyl, 100 mcg IM/IN. May repeat 50 mcg every 10 minutes titrated to pain, not to exceed 200 mcg.

Acute traumatic injuries, acute abdominal/flank pain, burn injuries, Cancer pain, Sickle Cell Crisis:

Fentanyl, 50 mcg slow IV/IO push over one (1) minute. May repeat every five (5) minutes titrated to pain, not to exceed 200 mcg IV/IO, **or**

Fentanyl, 100 mcg IM/IN. May repeat 50 mcg every 10 minutes titrated to pain, not to exceed 200 mcg.

Pacing, synchronized cardioversion:

Fentanyl, 50 mcg slow IV/IO over one (1) minute. May repeat in five (5) minutes titrated to pain, not to exceed 200 mcg.

Fentanyl, 100 mcg IN. May repeat 50 mcg every 10 minutes titrated to pain, not to exceed 200 mcg.

Reference #s 3050, 4060, 4080, 5010, 7010, 7020, 11020, 13030, 14070, 14090, 14100, 14240

Fentanyl - Pediatric (ALS)

Fentanyl, 0.5 mcg/kg slow IV/IO over one (1) minute. May repeat in five (5) minutes titrated to pain, not to exceed 100 mcg.

Fentanyl, 1 mcg/kg IM/IN, may repeat every 10 minutes titrated to pain not to exceed 200 mcg.

Reference #s 3050, 4080, 5010, 7010, 7020, 13030, 14180, 14190, 14240

Glucose - Oral - Adult (BLS, LALS, ALS)

Adult with blood glucose less than 80 mg/dL:

Glucose - Oral, one (1) tube for patients with an intact gag reflex and hypoglycemia.

Reference #s 7010, 7020, 13020, 14060, 14080, 14230

Glucose - Oral - Pediatric (BLS, LALS, ALS)

Hypoglycemia - Neonates (0 - 4 weeks) with blood glucose less than 35 mg/dL or pediatric patients (more than 4 weeks) with glucose less than 60 mg/dL:

Glucose - Oral, one (1) tube for patients with an intact gag reflex and hypoglycemia.

Reference #s 7010, 7020, 14170, 14160

Glucagon - Adult (LALS, ALS)

Glucagon, 1 mg IM/SC/IN, if unable to establish IV. May administer one (1) time only.

Beta blocker Poisoning (base hospital order only):

Glucagon, 1 mg IV/IO

Reference #s 4060, 4080, 7010, 7020, 13010, 13030, 14060

Glucagon - Pediatric (LALS, ALS)

Hypoglycemia, if unable to establish IV:

Glucagon, 0.03 mg/kg IM/IN, if unable to start an IV. May be repeated one (1) time after 20 minutes for a combined maximum dose of 1 mg.

Reference #s 7010, 7020, 13030, 14160, 14170

Beta blocker poisoning (base hospital order only):

Glucagon, 0.03 mg/kg IV/IO

Reference #'s 4060, 4080, 7010, 7020, 13010

Ipratropium Bromide (Atrovent) Inhalation Solution use with Albuterol Adult (ALS)

Atrovent, 0.5 mg nebulized. Administer one (1) dose only.

Reference #s 7010, 7020, 14010, 14070

Ipratropium Bromide (Atrovent) Metered-Dose Inhaler (MDI) use with Albuterol Adult (ALS - Specialty Programs Only)

When used in combination with Albuterol MDI use Albuterol MDI dosing.

Reference #s 4060, 4080, 7010, 7020, 14010, 14070

Ipratropium Bromide (Atrovent) Inhalation Solution use with Albuterol - Pediatric (ALS)

1 day to 12 months Atrovent, 0.25 mg nebulized. Administer one (1) dose only.
 1 year to 14 years Atrovent, 0.5 mg nebulized. Administer one (1) dose only.

Reference #s 7010, 7020, 14120, 14140, 14190

Ipratropium Bromide (Atrovent) Metered-Dose Inhaler (MDI) use with Albuterol - Pediatric (ALS - Specialty Programs Only)


When used in combination with Albuterol MDI use Albuterol MDI dosing.

Reference #s 4060, 4080, 7010, 7020, 14120, 14140, 14190

Ketamine - Adult (ALS)

Acute traumatic injury, acute abdominal/flank pain, burn injuries, cancer related pain and sickle cell crisis:
 Ketamine, 0.3 mg/kg to a max of 30 mg in a 50 - 100 ml of NS via IV over five (5) minutes. May repeat one (1) time, after 15 minutes, if pain score remains at five (5) or higher. Do not administer IVP, IO, IM, or IN.

This is the official pain scale to be used in patient assessment and documented on the PCR.



Reference #s 7010, 7020, 14100

Lidocaine - Adult (ALS)

VT (pulseless)/VF:

Initial Dose: Lidocaine, 1.5 mg/kg IV/IO

For refractory *VT (pulseless)/VF*, may administer an additional 0.75 mg/kg IV/IO, repeat one (1) time in five (5) to 10 minutes; maximum total dose of 3 mg/kg.

V-Tach, Wide Complex Tachycardia - with Pulses:

Lidocaine, 1.5 mg/kg slow IV/IO

May administer an additional 0.75 mg/kg slow IV/IO; maximum total dose of 3 mg/kg.

Reference #s 4060, 5010, 7010, 7020, 8010, 11020, 14040, 14050, 14090

Lidocaine - Pediatric (ALS)*Cardiac Arrest:*

1 day to 8 years Lidocaine, 1.0 mg/kg IV/IO

9 to 14 years Lidocaine, 1.0 mg/kg IV/IO

May repeat Lidocaine at 0.5 mg/kg after five (5) minutes; maximum total dose of 3 mg/kg.

*Reference #s 5010, 7010, 7020, 14150***Lidocaine 2% (Intravenous Solution) - Pediatric and Adult (ALS)***Pain associated with IO infusion:*

Lidocaine, 0.5 mg/kg slow IO push over two (2) minutes, not to exceed 40 mg total.

*Reference #s 5010, 7010, 7020, 11020***Magnesium Sulfate (ALS)***Polymorphic Ventricular Tachycardia:*

Magnesium Sulfate, 2 gm IV/IO bolus over five (5) minutes for polymorphic VT if prolonged QT is observed during sinus rhythm post-cardioversion.

Eclampsia (Seizure/Tonic/Clonic Activity):

Magnesium Sulfate, 4 gm IV/IO slow IV push over three (3) to four (4) minutes.

Magnesium Sulfate, 10 mg/min IV/IO drip to prevent continued seizures.

*Reference #s 5010, 7010, 7020, 8010, 14210**Severe Asthma/Respiratory Distress (ALS) (base hospital order only):*

Magnesium Sulfate, 2 gm slow IV drip over 20 minutes. Do not repeat.

*Reference# 14010***Magnesium Sulfate - Pediatric (ALS)***Severe Asthma/Respiratory Distress (base hospital order only):*

Magnesium Sulfate, 50 mg/kg slow IV drip over 20 minutes. Do not exceed the adult dosage of 2 gm total. Do not repeat.

*Reference # 14120***Midazolam (Versed) - Adult (ALS)***Behavioral Emergencies, with suspected excited delirium:*Midazolam, 2.5 mg IV/IO/IN. May repeat in five (5) minutes, **or**

Midazolam, 5 mg IM. May repeat in 10 minutes.

Maximum of three (3) doses using any combination of IV/IO/IM/IN may be administered. Contact base hospital for additional orders and to discuss further treatment options.

Reference # 14110

Seizure:

Midazolam, 2.5 mg IV/IO/IN. May repeat in five (5) minutes for continued seizure activity, **or**

Midazolam, 5 mg IM. May repeat in 10 minutes for continued seizure activity.

Assess patient for medication related reduced respiratory rate or hypotension.

Maximum of three (3) doses using any combination of IV/IO/IM/IN may be administered for continued seizure activity. Contact base hospital for additional orders and to discuss further treatment options.

Pacing, synchronized cardioversion:

Midazolam, 2 mg slow IV/IO push or IN

CPAP:

Midazolam, 1 mg slow IV/IO push may be administered one (1) time for anxiety related to application of CPAP. Contact base hospital for additional orders.

Reference #s 4060, 4080, 7010, 7020, 11020, 13020, 14060, 14210

Midazolam (Versed) - Pediatric (ALS)**Seizures:**

Midazolam, 0.1 mg/kg IV/IO with maximum dose 2.5 mg. May repeat Midazolam in five (5) minutes, **or**

Midazolam, 0.2 mg/kg IM/IN with maximum dose of 5 mg. May repeat Midazolam in 10 minutes for continued seizure.

Assess patient for medication related reduced respiratory rate or hypotension.

Maximum of three (3) doses using any combination of IV/IO/IM/IN may be administered for continued seizure activity. Contact base hospital for additional orders and to discuss further treatment options.

Reference #s 7010, 7020, 14170

Naloxone (Narcan) - Adult (BLS)**For resolution of respiratory depression related to suspected opiate overdose:**

Naloxone, 0.5 mg IM/IN, may repeat Naloxone 0.5 mg IM/IN every two (2) to three (3) minutes if needed.

For suspected Fentanyl overdose with respiratory depression:

Consider a loading dose of 4 mg IN Naloxone. If no signs of respiratory improvement, consider Naloxone 0.5 mg IM/IN every two (2) to three (3) minutes if needed.

Do not exceed 10 mg of Naloxone total regardless of route administered.

Reference #s 7010, 7020, 8030, 14060

Naloxone (Narcan) - Adult (LALS, ALS)

For resolution of respiratory depression related to suspected opiate overdose:

Naloxone, 0.5 mg IV/IO/IM/IN, may repeat Naloxone 0.5 mg IV/IO/IM/IN every two (2) to three (3) minutes if needed.

For suspected Fentanyl overdose with respiratory depression:

Consider a loading dose of 4 mg IN Naloxone. If no signs of respiratory improvement, consider Naloxone 0.5 mg IV/IO/IM/IN every two (2) to three (3) minutes if needed.

Do not exceed 10 mg of Naloxone total regardless of route administered.

Reference #s 4080, 7010, 7020, 14060

Naloxone (Narcan) - Pediatric (BLS)

For resolution of respiratory depression related to suspected opiate overdose:

1 day to 8 years	Naloxone, 0.1 mg/kg IM/IN (do not exceed the adult dose of 0.5 mg per administration)
9 to 14 years	Naloxone, 0.5 mg IM/IN

May repeat every two (2) to three (3) minutes if needed. Do not exceed the adult dosage of 10 mg total IM/IN.

Reference #s 7010, 7020, 8030, 14150, 14160

Naloxone (Narcan) - Pediatric (LALS, ALS)

For resolution of respiratory depression related to suspected opiate overdose:

1 day to 8 years	Naloxone, 0.1 mg/kg IV/IO/IM/IN (do not exceed the adult dose of 0.5 mg per administration)
9 to 14 years	Naloxone, 0.5 mg IV/IO/IM/IN

May repeat every two (2) to three (3) minutes if needed. Do not exceed the adult dosage of 10 mg total IV/IO/IM/IN.

Reference #s 7010, 7020, 14150, 14160

Nitroglycerin (NTG) (LALS, ALS)

Nitroglycerin, 0.4 mg sublingual/transmucosal.

One (1) every three (3) minutes as needed. May be repeated as long as patient continues to have signs of adequate tissue perfusion. **If a Right Ventricular Infarction is suspected, the use of nitrates requires base hospital contact.**

Nitroglycerin Paste, 1 inch (1 gm) transdermal, may not repeat.

Nitroglycerin sublingual is the preferred route of administration for ACS. Nitro Paste is a one (1) time dose and intended for when sublingual cannot be easily administered (i.e., CPAP).

Nitroglycerin is contraindicated if there are signs of inadequate tissue perfusion or if sexual enhancement medications have been utilized within the past 48 hours.

Reference #s 4060, 4080, 7010, 7020, 14010, 14240

Ondansetron (Zofran) - Patients four (4) years old to Adult (ALS)*Nausea/Vomiting:*

Ondansetron, 4 mg slow IV/IO/ODT

All patients four (4) to eight (8) years old: May administer a total of 4 mgs of Ondansetron prior to base hospital contact.

All patients nine (9) and older: May administer Ondansetron 4 mg; may repeat two (2) times, at 10 minute intervals, for a total of 12 mgs prior to base hospital contact.

May be used as prophylactic treatment of nausea and vomiting associated with narcotic administration.

Reference #s 4080, 7010, 7020, 14090, 14180, 14220

Oxygen (non-intubated patient per appropriate delivery device)*General Administration (Hypoxia):*

Titrate Oxygen at lowest rate required to maintain SPO₂ at 94%. Do not administer supplemental oxygen for SPO₂ more than 95%.

Chronic Obstructive Pulmonary Disease (COPD):

Titrate Oxygen at lowest rate required to maintain SPO₂ at 90%. Do not administer supplemental oxygen for SPO₂ more than 91%.

Reference #s 12010, 13010, 13020, 13030, 13050, 14010, 14020, 14030, 14040, 14060, 14070, 14090, 14120, 14130, 14140, 14160, 14170, 14180, 14190, 14200, 14210, 14220, 14230, 14240

Sodium Bicarbonate - Adult (ALS)*Tricyclic Poisoning (base hospital order only):*

Sodium Bicarbonate, 1 mEq/kg IV/IO

Reference #s 5010, 7010, 7020, 13010

For cardiac arrest with suspected metabolic acidosis, hyperkalemia or tricyclic poisoning (base hospital order only):

Sodium Bicarbonate, 50 mEq IV/IO

Reference #'s 7010, 7020, 14050

Sodium Bicarbonate - Pediatric (ALS)*Tricyclic Poisoning (base hospital order only):*

Sodium Bicarbonate, 1 mEq/kg IV/IO

Reference #'s 7010, 7020, 13010

Tranexamic Acid (TXA) - Patients 15 years of age and older (ALS)

Signs of hemorrhagic shock meeting inclusion criteria:

Administer TXA 1 gm in 50 - 100 ml of NS via IV/IO over 10 minutes. Do not administer IVP as this will cause hypotension.

Reference #s 7010, 7020, 14090

APPENDIX I**Medications for self-administration or with deployment of the ChemPack.**

Medications listed below may be used only for the purposes referenced by the associated ICEMA Treatment Protocol. Any other use, route or dose other than those listed, must be ordered in consultation with the Base Hospital physician.

Atropine - Pediatric (BLS, AEMT-Auto-injector only with training, ALS)

Known nerve agent/organophosphate poisoning with deployment of the ChemPack using:

Two (2) or more mild symptoms: Administer the weight-based dose listed below as soon as an exposure is known or strongly suspected. If severe symptoms develop after the first dose, two (2) additional doses should be repeated in rapid succession 10 minutes after the first dose; do not administer more than three (3) doses. If profound anticholinergic effects occur in the absence of excessive bronchial secretions, further doses of atropine should be withheld.

One (1) or more severe symptoms: Immediately administer (3) three weight-based doses listed below in rapid succession.

Weight-based dosing:

Less than 6.8 kg (less than 15 lbs):	0.25 mg, IM using multi-dose vial
6.8 to 18 kg (15 to 40 lbs):	0.5 mg, IM using AtroPen auto-injector
18 to 41 kg (40 to 90 lbs):	1 mg, IM using AtroPen auto-injector
More than 41 kg (more than 90 lbs):	2 mg, IM using multi-dose vial

Symptoms of insecticide or nerve agent poisoning, as provided by manufacturer in the AtroPen product labeling, to guide therapy:

Mild symptoms: Blurred vision, bradycardia, breathing difficulties, chest tightness, coughing, drooling, miosis, muscular twitching, nausea, runny nose, salivation increased, stomach cramps, tachycardia, teary eyes, tremor, vomiting, or wheezing.

Severe symptoms: Breathing difficulties (severe), confused/strange behavior, defecation (involuntary), muscular twitching/generalized weakness (severe), respiratory secretions (severe), seizure, unconsciousness, urination (involuntary).

NOTE: Infants may become drowsy or unconscious with muscle floppiness as opposed to muscle twitching.

Reference #s 11010, 13010, 13040

Diazepam (Valium) - Adult (ALS)

For seizures associated with nerve agent/organophosphate exposure ONLY with the deployment of the ChemPack:

Diazepam 10 mg (5 mg/ml) auto-injector IM (if IV is unavailable), **or**
Diazepam 2.5 mg IV

Reference # 13040

Diazepam (Valium) - Pediatric (ALS)

For seizures associated with nerve agent/organophosphate exposure ONLY with the deployment of the ChemPack:

Diazepam 0.05 mg/kg IV

Reference # 13040

Nerve Agent Antidote Kit (NAAK)/Mark I or DuoDote (containing Atropine/Pralidoxime Chloride for self-administration or with deployment of the ChemPack) - Adult

Nerve agent exposure with associated symptoms:

One (1) NAAK auto-injector IM into outer thigh. May repeat up to two (2) times every 10 to 15 minutes if symptoms persist.

Reference #s 7010, 7020, 13010, 13040



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 14110R1

Effective Date: 10/01/21

Supersedes: 03/01/20

Page 1 of 2

PSYCHIATRIC/BEHAVIORAL EMERGENCIES - ADULT

I. PURPOSE

To provide timely and appropriate treatment for patients that are violent, potentially violent, or who may harm themselves or others, including the potential use of restraints in the field or during transport.

II. FIELD ASSESSMENT/TREATMENT INDICATORS

Symptoms of dangerous agitation, confusion and hallucinations, erratic behavior, profuse diaphoresis, elevated vital signs, hyperthermia, unexplained strength and endurance, and behaviors that include clothing shedding, shouting, and extreme thrashing when restrained.

This policy is not intended to negate the need for law enforcement personnel to use appropriate restraint equipment to establish scene-management control. Restraints should be applied by law enforcement whenever possible. If applied, an officer is required to remain available at the scene or during transport to remove or adjust the restraints for patient safety per ICEMA Reference #6060 - Patient Restraints.

III. BLS INTERVENTIONS

- Approach patient in a calm and cautious manner.
- Ensure patent airway, obtain oxygen saturation and apply oxygen as needed.
- Restraint equipment must be either padded leather restraints or soft restraints (e.g., posey, velcro or seat-belt type).
- Apply four (4) point restraints as clinically indicated. Transport of a restrained patient should be in low to high Fowlers position. Never transport a patient in a prone position while restrained. Transport of a patient supine, while restrained, can affect respiratory function and constant monitoring of respiratory status is required.
- Perform cooling measures as clinically indicated.
- If suspected hypoglycemia, obtain a blood glucose.

IV. LIMITED ALS (LALS) INTERVENTIONS

- Perform activities identified in the BLS Interventions.

V. ALS INTERVENTIONS

- Perform activities identified in the BLS and LALS Interventions.
- If patient meets criteria for potentially fatal and dangerous agitation, administer Midazolam per ICEMA Reference #11010 - Medication - Standard Orders. Do not delay administration of Midazolam due to lack of vascular access as IM or IN is preferred in this circumstance. May repeat one (1) time using same method as first administered.

- Place on cardiac monitor. Continuous monitoring of a patient after administration of Midazolam is required.
- Obtain capnography, monitor waveform and numerical value. Apnea can be the result of the use of Midazolam and other medications.
- Once conditions are safe, establish IV.
- Base hospital may order:
 - For potentially fatal and dangerous agitation and suspected metabolic acidosis/hyperkalemia administer Sodium Bicarbonate per ICEMA Reference #11010 - Medication - Standard Orders.

VI. REFERENCES

<u>Number</u>	<u>Name</u>
6060	Patient Restraints
11010	Medication - Standard Orders



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. **4060R12**
Effective Date: 10/01/201
Supersedes: ~~03/01/2010~~10/01/20
Page 1 of 3

FIRELINE EMT-P

I. PURPOSE

To provide guidance and medical oversight for an ICEMA paramedic (EMT-P) deployed to function as a fireline paramedic (FEMP).

This protocol is for use by authorized FEMPs during fire suppression activities and treatment of fire suppression personnel only.

II. REQUIREMENTS

- Must be a currently licensed EMT-P in California.
- Must be currently accredited EMT-P in the ICEMA region.
- Must be currently employed by an ICEMA approved ALS provider.
- The FEMP will follow FIRESCOPE FEMP ICS 223-11 Position Manual and all other ICS protocols.
- The FEMP will check in and obtain briefing from the Logistics Section Chief or the Medical Unit Leader, if established. Briefing will include current incident situation, anticipated medical needs, and local emergency medical system orientation.
- The FEMP will provide emergency medical treatment to personnel operating on the fireline.
- The FEMP will follow ICEMA prior to contact protocols if unable to contact the assigned base hospital.
- The FEMP may not perform skills outside of the ICEMA scope of practice.

III. PROCEDURE

- The EMS provider will notify ICEMA of the deployment of the FEMP to an incident. Use the Fireline Paramedic (FEMP) Deployment Notification form, which is on the ICEMA website at ICEMA.net.
- The FEMP will carry inventory in the advanced life support (ALS) pack as per the below inventory list (see Section IV. Fireline EMT-P (ALS) Pack Inventory. Inventory will be supplied and maintained by the employing ALS provider. Additional items for restock should also be maintained and secured in a vehicle or in the Medical Unit trailer.
- Incident Medical Units may not have the capability of resupplying controlled substances (narcotics). EMS providers should stock sufficient quantities of medical supplies and medications, especially controlled substance medications, to assure adequate supplies and medications.
- Narcotics must be under double lock and maintained on the FEMP person or secured in his/her vehicle at all times as per the ICEMA Drug and Equipment List.

- FEMP may carry an inventory of controlled substances (i.e., Fentanyl, Ketamine and Midazolam) if authorized by the employing ALS provider's Medical Director. The authorizing Medical Director is responsible to assure full compliance with all federal and state laws relating to purchase, storage and transportation of controlled substances. Only controlled substances approved for use in the ICEMA region may be carried and their use must be in accordance with current ICEMA patient care protocols.
- Radio communication failure protocols will not be used. Prior to base hospital contact protocols will be followed. If further treatment is needed, radio contact with the base hospital should be established as soon as possible.
- Documentation of patient care must follow ICEMA protocol utilizing the electronic patient care report (ePCR), if available, or a paper O1A form. All PCRs will be reviewed by the ALS provider and ICEMA for quality improvement (QI) purposes.
- A FEMP will be paired with a fireline EMT (FEMT) or another FEMP who will assist with basic life support (BLS) treatment and supplies.

IV. FIRELINE EMT-P (ALS) PACK INVENTORY

Minimum Requirements: The weight of the pack will dictate if the EMT-P chooses to carry additional ALS supplies.

MEDICATIONS/SOLUTIONS

Medications/Solutions	ALS
Albuterol Solution 2.5 mg Handheld Nebulizer or Multidose Inhaler	4
Atropine Sulfate 1 mg	2
Ipratropium Bromide Solution 0.5 mg Handheld Nebulizer or Multidose Inhaler	4
Lidocaine 100 mg IV pre-load	2
Aspirin 80 mg chewable	1 bottle
Dextrose 10%/250 ml (D10W 25 gm) IV/IO Bolus	1
Diphenhydramine 50 mg	4
Epinephrine 1: 10,000 1 mg	2
Epinephrine 1: 1000 1 mg	4
Glucagon 1 mg	1
<u>Naloxone 2 mg</u>	<u>2</u>
Nitroglycerin - Spray 0.4 metered dose and/or tablets (tablets to be discarded 90 days after opening)	1 (equivalent of 10 patient doses)
Saline 0.9% IV 1000 ml may be divided in two 500 ml bags or four 250 ml bags.	
Tranexamic Acid (TXA) 1 gm	1

CONTROLLED SUBSTANCE MEDICATIONS

Controlled Substance Medications MUST BE DOUBLED LOCKED	ALS
Midazolam	20 mg
Fentanyl (amount determined by the medical director)	200 - 400 mcg
Ketamine	120 - 1000 mg

ALS AIRWAY EQUIPMENT

Airway Equipment	ALS
Endotracheal Tubes - 6.0, 7.0 and/or 7.5 cuffed with stylet	1 each
Laryngeal blades - #0, #1, #2, #3, #4 curved and/or straight	1 each
Laryngoscope handle with batteries - or 2 disposable handles	1 each
ET Tube holder	1
End Tidal CO2 Detector	1
Needle Cricothyrotomy Kit	1
Needle Thoracostomy Kit	1

IV/MEDICATION ADMINISTRATION SUPPLIES

IV/Medication Administration Supplies	ALS
IV administration set macro drip	2
Venaguard	2
Alcohol preps	6
Betadine swabs	4
Tourniquet	2
Razor	1
Tape	1
IV catheters - 14, 16, 18 and 20 gauge	2
10 cc syringe	2
1 cc TB syringe	2
18 gauge needle	4
25 gauge needle	2

MISCELLANEOUS EQUIPMENT

Miscellaneous	ALS
Sharps container	1
Narcotic storage per protocol	
FEMP pack inventory sheet	1
Patient care report or ePCR (Toughbook)	
AMA forms	3

Equipment	ALS
Compact AED or compact monitor defibrillator combination	
Appropriate cardiac pads	
Pulse oximetry (optional)	
Glucometer, test strips and lancets	4

The BLS pack and supplies will be carried by the FEMT or accompanying FEMP. Personal items and supplies cannot be carried in either the ALS pack or the BLS pack.



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. **6020R1**
Effective Date: ~~03/01/2010/01/21~~
Supersedes: ~~06/18/1903/01/20~~
Page 1 of 1

RESPONSIBILITY FOR PATIENT MANAGEMENT

I. PURPOSE

To define the responsibility for patient care management in the prehospital setting. Within the ICEMA region, in the event both public and private emergency medical services (EMS) field personnel arrive on the scene with the same qualifications, patient care management responsibility will rest with the first to arrive.

II. PROCEDURE

- An advanced emergency medical technician (AEMT) or paramedic (EMT-P) may transfer patient management responsibility to an emergency medical technician (EMT) for transportation, ~~without base hospital direction~~, only under the following conditions:
 - The patient is stable for transport and no ALS measures have been initiated.
 - When operating under ICEMA Reference #8030 - Transport of Patients (BLS). When the patient does not meet criteria for base hospital contact and has not received advanced life support (ALS) care.
 - When operating under ICEMA Reference #80980 - Medical Response to a Multiple Casualty Incident.
 - When operating under ICEMA Reference #6050 - Local Medical Emergency.
- The base hospital should be contacted if at any time transfer of patient management responsibility is in question or for any patient not meeting the above criteria.
- ~~In the event of radio communication failure, a limited ALS (LALS) or ALS unit may not transfer patient management responsibility to an EMT for transportation.~~

III. REFERENCES

<u>Number</u>	<u>Name</u>
6050	Local Medical Emergency
8030	Transport of Patients (BLS)
8080	Medical Response to a Multiple Casualty Incident



**INLAND COUNTIES
EMERGENCY MEDICAL AGENCY
POLICY AND PROTOCOL MANUAL**

Reference No. 7010R23
 Effective Date: ~~06/01/21~~ 10/01/21
 Supersedes: ~~10/01/2006~~ 01/21
 Page 1 of 6

STANDARD DRUG AND EQUIPMENT LIST - BLS/LALS/ALS

Each ambulance and first responder unit shall be equipped with the following functional equipment and supplies. **This list represents mandatory items with minimum quantities** excluding narcotics, which must be kept within the range indicated. All expiration dates must be current. All packaging of drugs or equipment must be intact. No open products or torn packaging may be used.

All ALS (transport and non-transport) and BLS transport vehicles shall be inspected annually.

MEDICATIONS/SOLUTIONS

Exchanged Medications/Solutions	BLS	LALS	ALS Non-Transport	ALS Transport
Adenosine (Adenocard) 6 mg			1	1
Adenosine (Adenocard) 12 mg			2	2
Albuterol Aerosolized Solution (Proventil) - unit dose 2.5 mg		4 doses	4 doses	4 doses
Albuterol MDI with spacer		1 SPECIALTY PROGRAMS ONLY	1 SPECIALTY PROGRAMS ONLY	1 SPECIALTY PROGRAMS ONLY
Aspirin, chewable - 81 mg tablet		2	1 bottle	1 bottle
Atropine 1 mg preload			2	2
Calcium Chloride 1 gm preload			1	1
Dextrose 10% in 250 ml Water (D10W)		2	2	2
Diphenhydramine (Benadryl) 50 mg			1	1
Epinephrine 1 mg/ml 1 mg		2	2	2
Epinephrine 0.1 mg/ml 1 mg preload			4	4
Glucagon 1 mg		1	1	1
Glucose paste	1 tube	1 tube	1 tube	1 tube
Ipratropium Bromide Inhalation Solution (Atrovent) unit dose 0.5 mg			4	4
Irrigating Saline and/or Sterile Water (1000 cc)	2	1	1	2
Lidocaine 100 mg			3	3
Lidocaine 2% Intravenous solution			1	1
Magnesium Sulfate 10 gm			1	1
Naloxone (Narcan) 2 mg preload	2	2	2	2
Nitroglycerine (NTG) - Spray 0.4 mg metered dose and/or tablets (tablets to be discarded 90 days after opening)		2	1	2
Nitroglycerine Paste 2% - 1 gm packets, or Nitroglycerine Paste 2% - 30 gm tube, or Nitroglycerine Paste 2% - 60 gm tube				2 1 1
Normal Saline for Injection (10 cc)		2	2	2
Normal Saline 100 cc			1	2
Normal Saline 250 cc			1	1
Normal Saline 500 ml and/or 1000 ml		2000 ml	3000 ml	6000 ml
Ondansetron (Zofran) 4 mg Oral Disintegrating Tablets (ODT)			4	4
Ondansetron (Zofran) 4 mg IM/IV			4	4
Sodium Bicarbonate 50 mEq preload			2	2
Tranexamic Acid (TXA) 1 gm			2	2

Non-Exchange Controlled Substance Medications MUST BE DOUBLE LOCKED	BLS	LALS	ALS Non- Transport	ALS Transport
Fentanyl			200-400 mcg	200-400 mcg
Midazolam			20-40 mg	20-40 mg
Ketamine			120-1000 mg	120-1000 mg

AIRWAY/SUCTION EQUIPMENT

Exchanged Airway/Suction Equipment	BLS	LALS	ALS Non- Transport	ALS Transport
CPAP circuits - all manufacture's available sizes			1 each	2 each
End-tidal CO2 device - Pediatric and Adult (may be integrated into bag)			1 each	1 each
Endotracheal Tubes cuffed - 6.0 and/or 6.5, 7.0 and/or 7.5 and 8.0 and/or 8.5 with stylet			2 each	2 each
ET Tube holders - adult		1 each	1 each	2 each
Mask - Adult & Pediatric non-rebreather oxygen mask	2 each	2 each	2 each	2 each
Mask - Infant Simple Mask	1	1	1	1
Nasal cannulas - pediatric and adult	2 each	2 each	2 each	2 each
Naso/Orogastric feeding tubes - 5fr or 6fr, and 8fr			1 each	1 each
Naso/Orogastric tubes - 10fr or 12fr, 14fr, 16fr or 18fr			1 each	1 each
Nasopharyngeal Airways - (infant, child, and adult)	1 each	1 each	1 each	1 each
Needle Cricothyrotomy Device - Pediatric and adult or Needles for procedure 10, 12, 14 and/or 16 gauge			1 each 2 each	1 each 2 each
One way flutter valve with adapter or equivalent			1	1
Oropharyngeal Airways - (infant, child, and adult)	1 each	1 each	1 each	1 each
Rigid tonsil tip suction	1		1	1
Small volume nebulizer with universal cuff adaptor		2	2	2
Suction Canister	1		1	1
Suction catheters - 6fr, 8fr or 10fr, 12fr or 14fr	1 each		1 each	1 each
Ventilation Bags - Infant 250 ml Pediatric 500 ml (or equivalent) Adult	1 1 1	1 1 1	1 1 1	1 1 1
Water soluble lubricating jelly		1	1	1

Non-Exchange Airway/Suction Equipment	BLS	LALS	ALS Non-Transport	ALS Transport
Ambulance oxygen source -10 L / min for 20 minutes	1			1
CPAP - (must be capable of titrating pressure between 2 and 15 cm H ₂ O)			1	1
Flashlight/penlight	1	1	1	1
Laryngoscope blades - #0, #1, #2, #3, #4 curved and/or straight			1 each	1 each
Laryngoscope handle with batteries - or 2 disposable handles			1	1
Magill Forceps - Pediatric and Adult			1 each	1 each
Manual powered suction device		1		
Portable oxygen with regulator - 10 L /min for 20 minutes	1	1	1	1
Portable suction device (battery operated)	1		1	1
Pulse Oximetry device	(SEE OPTIONAL EQUIPMENT SECTION, PG. 5)	1	1	1
Stethoscope	1	1	1	1
Wall mount suction device	1 (BLS TRANSPORT ONLY)			1

IV/NEEDLES/SYRINGES/MONITORING EQUIPMENT

Exchanged IV/Needles/Syringes/Monitor Equipment	BLS	LALS	ALS Non-Transport	ALS Transport
Conductive medium or Pacer/Defibrillation pads			2 each	2 each
Disposable Tourniquets		2	2	2
ECG electrodes			20	20
EZ-IO Driver			1 each	1 each
EZ-IO Needles: 25 mm 45 mm			2 each 1 each	2 each 1 each
Glucose monitoring device with compatible strips and OSHA approved single use lancets	1	1	1	1
3-way stopcock with extension tubing			2	2
IV Catheters - sizes 14, 16, 18, 20, 22, 24		2 each	2 each	2 each
Macro drip Administration Set		3	3	3
Micro drip Administration Set (60 drops / cc)		1	1	2
Mucosal Atomizer Device (MAD) for nasal administration of medication	2	2	2	4
Pressure Infusion Bag (disposable)		1	1	1
Razors		1	2	2
Safety Needles - 20 or 21gauge and 23 or 25 gauge	2 each	2 each	2 each	2 each
Saline Lock Large Bore Tubing Needleless		2	2	2
Sterile IV dressing		2	2	2
Syringes w/wo safety needles - 1 cc, 3 cc, 10 cc catheter tip		2 each		

Exchanged IV/Needles/Syringes/Monitor Equipment	BLS	LALS	ALS Non-Transport	ALS Transport
Syringes w/wo safety needles - 1 cc, 3 cc, 10 cc, 20 cc, 60 cc catheter tip			2 each	2 each

Non-Exchange IV/Needles/Syringes/ Monitor Equipment	BLS	LALS	ALS Non-Transport	ALS Transport
12-lead ECG Monitor and Defibrillator with TCP and printout			1	1
Blood pressure cuff - large adult or thigh cuff, adult, child and infant (one of each size)	1	1	1	1
Capnography monitor and supplies, may be integrated in the cardiac monitor			1	1
Needle disposal system (OSHA approved)	1	1	1	1
Thermometer - Mercury Free with covers	1	1	1	1

OPTIONAL EQUIPMENT/MEDICATIONS

Non-Exchange Optional Equipment/ Medications	BLS	LALS	ALS Non-Transport	ALS Transport
AED/defib pads - Adult (1), Pediatric (1)	1 each	1 each		
Automatic CPR device (FDA approved)	1	1	1	1
Automatic transport ventilator (Specialty Program Only - ICEMA approved device)			1	1
Backboard padding	1	1	1	1
Buretrol			1	1
Chemistry profile tubes			3	3
CPAP – (must be capable of titrating pressure between 2 and 15 cm H₂O)			4	4
Nerve Agent Antidote Kit (NAAK) - DuoDote or Mark I	3	3	3	3
EMS Tourniquet	1		1	1
Gum Elastic intubation stylet			2	2
Hemostatic Dressings *	1	1	1	1
IO Needles - Manual, Adult and Pediatric, Optional		Pediatric sizes only or EZ-IO needles and drivers	1 each	1 each
IV infusion pump			1	1
IV warming device		1	1	1
Manual IV Flow Rate Control Device			1	1
Manual powered suction device	1	1	1	1
Multi-lumen peripheral catheter			2	2
Needle Thoracostomy Kit (prepackaged)			2	2
<u>Naloxone (Narcan) Nasal Spray 4 mg</u>	<u>2</u>	<u>2</u>	<u>2</u>	<u>2</u>
Pulse Oximetry device	1			
Translaryngeal Jet Ventilation Device			1	1
Vacutainer			1	1

* Hemostatic Dressings

- Quick Clot, Z-Medica
 - Quick Clot, Combat Gauze LE
 - Quick Clot, EMS Rolled Gauze, 4x4 Dressing, TraumaPad
- Celox

- Celox Gauze, Z-Fold Hemostatic Gauze
- Celox Rapid, Hemostatic Z-Fold Gauze
- HemCon ChitoFlex Pro Dressing

NOTE:

- The above products are “packaged” in various forms (i.e., Z-fold, rolled gauze, trauma pads, 4”x4”pads) and are authorized provided they are comprised of the approved product.
- Hemostatic Celox Granules, or granules delivered in an applicator, are not authorized.

DRESSING MATERIALS/OTHER EQUIPMENT/SUPPLIES

Exchanged Dressing Materials/Other Equipment/Supplies	BLS	LALS	ALS Non-Transport	ALS Transport
Adhesive tape - 1 inch	2	2	2	2
Air occlusive dressing	1	1	1	1
Ankle and wrist restraints, soft ties acceptable	1		1	1
Antiseptic swabs/wipes	10	10	10	10
Bedpan or fracture pan	1 (BLS TRANSPORT UNITS ONLY)			1
Urinal	1 (BLS TRANSPORT UNITS ONLY)			1
Cervical Collars - Rigid Pediatric and Adult all sizes or	2 each	2 each	2 each	2 each
Cervical Collars - Adjustable Adult and Pediatric	2 each	2 each	2 each	2 each
Cold Packs	2	2	2	2
Emesis basin or disposable bags and covered waste container	1	1	1	1
Head immobilization device	2	2	2	2
OB Kit	1	1	1	1
Pneumatic or rigid splints capable of splinting all extremities	4	2	2	4
Provodine/Iodine swabs/wipes or antiseptic equivalent		4	10	10
Roller bandages - 4 inch	6	3	3	6
Sterile bandage compress or equivalent	6	2	2	6
Sterile gauze pads - 4x4 inch	4	4	4	4
Sterile sheet for Burns	2	2	2	2
Universal dressing 10x30 inches	2	2	2	2

Non-Exchange Dressing Materials/Other Equipment/Supplies	BLS	LALS	ALS Non-Transport	ALS Transport
800 MHz Radio		1	1	1
Ambulance gurney	1 (BLS TRANSPORT UNITS ONLY)			1
Bandage shears	1	1	1	1
Blood Borne Pathogen Protective Equipment - (nonporous gloves, goggles face masks and gowns meeting OSHA	2	1	2	2

Non-Exchange Dressing Materials/Other Equipment/Supplies	BLS	LALS	ALS Non-Transport	ALS Transport
Standards)				
Pediatric Emergency Measuring Tape (Broselow, etc.)		1	1	1
Drinkable water in secured plastic container or equivalent	1 gallon			1 gallon
Long board with restraint straps	1	1	1	1
Pediatric immobilization board	1	1	1	1
Pillow, pillow case, sheets and blanket	1 set (BLS TRANSPORT UNITS ONLY)			1 set
Short extrication device	1	1	1	1
Straps to secure patient to gurney	1 set (BLS TRANSPORT UNITS ONLY)			1 set
Traction splint	1	1	1	1
Triage Tags - ICEMA approved	20	20	20	20



**INLAND COUNTIES
EMERGENCY MEDICAL AGENCY
POLICY AND PROTOCOL MANUAL**

Reference No. 7020R42
Effective Date: 10/01/201
Supersedes: ~~03/01/2010/01/20~~
Page 1 of 4

STANDARD DRUG AND EQUIPMENT LIST - EMS AIRCRAFT

Each Aircraft shall be equipped with the following functional equipment and supplies. This list represents mandatory items with minimum quantities, to exclude narcotics, which must be kept within the range indicated. All expiration dates must be current. All packaging of drugs or equipment must be intact. No open products or torn packaging may be used.

MEDICATIONS/SOLUTIONS	AMOUNT
Adenosine (Adenocard) 6 mg	1
Adenosine (Adenocard) 12 mg	2
Albuterol Aerosolized Solution (Proventil) - unit dose 2.5 mg	3 doses
Aspirin, chewable - 81 mg tablet	1 bottle
Atropine 1 mg preload	2
Calcium Chloride 1 gm preload	1
Dextrose 10% in 250 ml Water (D10W) *	2
Diphenhydramine (Benadryl) 50 mg	1
Epinephrine 1 mg/ml 1 mg	2
Epinephrine 0.1 mg/ml 1mg preload	3
Glucagon 1 mg	1
Glucopaste	1 tube
Ipratropium Bromide Inhalation Solution (Atrovent) unit dose 0.5 mg	3
Lidocaine 100 mg	3
Lidocaine 2% Intravenous solution	1
Magnesium Sulfate 10 gms	1
Naloxone (Narcan) 2 mg preload	2
Nitroglycerin (NTG) - Spray 0.4 mg metered dose and/or tablets (tablets to be discarded 90 days after opening.)	1
Normal Saline for Injection (10 cc)	2
Normal Saline 250 ml	1
Normal Saline 500 ml and/or 1000 ml	2000 ml
Ondansetron (Zofran) 4 mg Oral Disintegrating Tablets (ODT)	4
Ondansetron (Zofran) 4 mg IM/ IV	4
Sodium Bicarbonate 50 mEq preload	2
Tranexamic Acid (TXA) 1 gm	2

CONTROLLED SUBSTANCE MEDICATIONS-MUST BE DOUBLE LOCKED	AMOUNT
Fentanyl	200-400 mcg
Midazolam	20-40 mg
Ketamine	120-1000 mg

AIRWAY/SUCTION EQUIPMENT	AMOUNT
Aircraft Oxygen source -10 L /min for 20 minutes	1
C-PAP circuits - all manufacture's available sizes	1 each
End-tidal CO2 device - pediatric and adult (may be integrated into bag)	1 each
Endotracheal Tubes cuffed - 6.0 and/or 6.5, 7.0 and/or 7.5 and 8.0 and/or 8.5 with stylet	2 each
ET Tube holders - adult	1 each
Flashlight/penlight	1
Laryngoscope handle with batteries - or 2 disposable handles	1
Laryngoscope blades - #0, #1, #2, #3, #4 curved and/or straight	1 each
Magill Forceps - Pediatric and Adult	1 each
Nasal Cannulas - infant, pediatric and adult	2 each
Naso/Orogastric tubes - 10fr or 12fr, 14fr, 16fr or 18fr	1 each

AIRWAY/SUCTION EQUIPMENT	AMOUNT
Naso/Orogastric feeding tubes - 5fr or 6fr, and 8fr	1 each
Nasopharyngeal Airways - infant, child, and adult	1 each
Needle Cricothyrotomy Device (Approved) - Pediatric and adult <i>or</i>	1 each
Needles for procedure 10, 12, 14 and/or 16 gauge	2 each
Non Re-Breather O ₂ Mask - Pediatric and Adult, Infant Simple Mask	2 each
One way flutter valve with adapter or equivalent	1
Oropharyngeal Airways - infant, child, and adult	1 each
Portable Oxygen with regulator - 10 L /min for 20 minutes	1
Portable suction device (battery operated) <i>and/or</i> Wall mount suction device	1 each
Pulse Oximetry device	1
Small volume nebulizer with universal cuff adaptor	1
Stethoscope	1
Suction catheters - 6fr, 8fr or 10fr, 12fr or 14fr	1 each
Ventilation Bags - Infant 250 ml, Pediatric 500 ml and Adult 1 L	1 each
Water soluble lubricating jelly	1
Ridged tonsil tip suction	1

IV/NEEDLES/SYRINGES/MONITORING EQUIPMENT	AMOUNT
12-Lead ECG Monitor and Defibrillator with TCP and printout	1
800 MHz Radio	1
Blood pressure cuff - large adult or thigh cuff, adult, child and infant	1 set
Capnography monitor and supplies, may be integrated in the cardiac monitor	1
Conductive medium <i>or</i> Adult and Pediatric Pacer/Defibrillation pads	2 each
ECG - Pediatric and Adult	20 patches
EZ IO Needles and Driver 25 mm <i>and</i>	2 each
45 mm	1 each
3-way stopcock with extension tubing	2
IO Needles - Manual, Adult and Pediatric, <u>Optional</u>	1 each
IV Catheters - sizes 14, 16, 18, 20, 22, 24	2 each
Glucose monitoring device	1
Macro drip Administration Set	3
Micro drip Administration Set (60 drops/ml)	1
Mucosal Atomizer Device (MAD) for nasal administration of medication	4
Needle disposal system (OSHA approved)	1
Pressure infusion bag	1
Safety Needles - 20 or 21 gauge and 23 or 25 gauge	2 each
Saline Lock	2
Syringes w/wo safety needles - 1 ml, 3 ml, 10 ml, 20 ml	2 each
Syringe - 60 ml catheter tip	2
Thermometer - Mercury free with covers	1

DRESSING MATERIALS/OTHER EQUIPMENT SUPPLIES	AMOUNT
Adhesive tape - 1 inch	2
Air occlusive dressing	1
Aircraft stretcher or litter system with approved FAA straps that allows for Axial Spinal Immobilization	1
Ankle and wrist restraints, soft ties acceptable	1
Antiseptic swabs/wipes	
Bandage shears	1
Blanket or sheet	2
Blood Borne Pathogen Protective Equipment - (nonporous gloves, goggles face masks and gowns meeting OSHA Standards)	2
Cervical Collars - Rigid Pediatric & Adult all sizes or Cervical Collars - Adjustable Adult and Pediatric	1 each 1 each
Emesis basin or disposable bags and covered waste container	1
Head immobilization device	1
OB Kit	1
Pediatric Emergency Measuring Tape (Broselow, etc.)	1
Pneumatic or rigid splints capable of splinting all extremities	4
Providine/Iodine swabs/wipes or antiseptic equivalent	
Roller bandages - 4 inch	3
Sterile bandage compress or equivalent	6
Sterile gauze pads - 4x4 inch	4
Sterile Sheet for Burns	2
Traction splint	1
Universal Dressing 10x30 inches	2

OPTIONAL EQUIPMENT/MEDICATIONS	Amount
Automatic ventilator (Approved)	1
Backboard padding	1
BLS AED/defib pads	1
Chemistry profile tubes	3
Nerve Agent Antidote Kit (NAAK) - DuoDote or Mark I	3
D5W in bag	1
Hemostatic Dressing *	1
IV infusion pump	1
IV warming device	1
Manual powered suction device	1
Medical Tourniquet	1
<u>Naloxone (Narcan) Nasal Spray 4 mg</u>	<u>2</u>
Needle Thoracostomy Kit (prepackaged)	2
Pediatric immobilization board	1
Translaryngeal Jet Ventilation Device	1
Vacutainer	1

* Hemostatic Dressings

- Quick Clot, Z-Medica
 - Quick Clot, Combat Gauze LE
 - Quick Clot, EMS Rolled Gauze, 4x4 Dressing, TraumaPad
- Celox
 - Celox Gauze, Z-Fold Hemostatic Gauze
 - Celox Rapid, Hemostatic Z-Fold Gauze

NOTE:

- The above products are “packaged” in various forms (i.e., Z-fold, rolled gauze, trauma pads, and 4”x4” pads) and are authorized provided they are comprised of the approved product.
- Hemostatic Celox Granules, or granules delivered in an applicator, are not authorized.



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. 8130R23

Effective Date: ~~12/10/2010/01/21~~

Supersedes: ~~12/02/2012/10/20~~

Page 1 of 3

ASSESS AND REFER: ~~EMERGENCY~~ RESPONSE PLAN (San Bernardino County Only)

I. PURPOSE

To establish standards for the identification of patients whose condition does not require transport by 9-1-1 emergency ambulance. All 9-1-1 calls for EMS will receive an appropriate response, timely assessment, and appropriate patient care. If it is determined that the patient is stable, and does not require emergency department services EMS field personnel will assess patient and provide an appropriate alternative recommendation.

II. POLICY

- ~~• This emergent policy is being implemented in accordance with the XBO COVID19 EMS Plan and will be used only in response to surge triggers defined in that plan in order to maintain continuity of EMS in San Bernardino County during a public health emergency.~~
- ~~• Destination decisions will be based on patient's condition or on patient, guardian, family or law enforcement requests.~~ If the patient's condition is stable and meets assess and refer criteria EMS field personnel will provide the patient the following recommendation:
 - "It appears that you do not require immediate care in the emergency department. You should seek care with your regular healthcare provider, urgent care or clinic. If symptoms worsen seek medical help or re-contact 9-1-1."

III. GENERAL CONSIDERATIONS

- Transport all patients requiring immediate medical attention to the closest most appropriate hospital.
- EMS should not require patients that are being released from the scene to sign AMA on the Patient Care Record.
- Provide instructions that if symptoms worsen, patient should go to the emergency department, contact their healthcare provider, or re-contact 9-1-1.
- If the patient or guardian refuses the referral, the patient will be transported to the closest most appropriate hospital.

IV. PARAMEDIC ASSESS AND REFER DECISION MAKING PRINCIPLES

- Does the patient, parent, or guardian have Decision Making Capacity?
- Is EMS field personnel concerned with the patient's current medical condition?
- How likely is the patient to successfully navigate the provided referral?

V. ASSESS AND REFER CRITERIA

- The patient must meet all of the following criteria:
 - Parent or guardian is on scene if the patient is under 18 years of age (unless legally emancipated).

- Has a Glasgow Coma Scale (GSC) of 15 or GCS is at patient's baseline.
- Exhibits no clinical evidence of:
 - Altered level of consciousness
 - Alcohol or drug ingestion that impairs decision making capacity
 - Abnormal or labored breathing or shortness of breath
 - Chest pain/discomfort of any kind
 - Hypoxia as indicated by low oxygen saturation
 - Significant tachycardia
 - Serious hemorrhage
- Exhibits evidence of Decision-Making Capacity sufficient to understand the nature of the medical condition as well as the risks and potential consequences of not seeking additional medical care from the provided recommendation.
- The patient would benefit from the provided recommendation.
- The patient is likely to successfully navigate the provided recommendation.
- If there is clinical evidence of a viral illness, the patient must meet all the following criteria: The COVID positive patient or person under investigation (PUI) must meet all of the following criteria:
 - Be stable.
 - Not under two (2) years of age, or over 65 years of age.
 - Does not have an underlying medical history.
- For the COVID positive patient or PUI, assess for a referral to stay home, self-isolate, and seek follow-up treatment with a physician.

VI. DOCUMENTATION REQUIREMENTS

- Physical exam.
- Treatment provided.
- Patient, parent, or guardian is alert, oriented, and acting appropriately for their age.
- Indications that there were no signs of significant impairment due to drugs, alcohol, organic causes, or mental illness.
- Any other observations that indicate that the patient, parent, or guardian has impaired Decision-Making Capacity.
- Recommendation/referrals shall be documented utilizing the following four (4) step process:
 - That a recommendation/referral was offered.
 - What the recommendation/referral was that EMS field personnel provided.
 - The patient's understanding of the recommendation/referral.

- The patient's plan based on the recommendation/referral of the EMS field personnel.
- The person(s), if any, who remained to look after the patient (the patient's "support system").
- The name of the interpreter utilized, if applicable.
- EMS field personnel will leave a referral card containing relevant community referral information with the patient.

~~NOTE: All assess and refer cases will undergo 100% CQI.~~



**INLAND COUNTIES
EMERGENCY MEDICAL AGENCY
POLICY AND PROTOCOL MANUAL**

Reference No. 11010R23
Effective Date: ~~06/15/21~~10/01/21
Supersedes: ~~06/01/21~~06/15/21
Page 1 of 13

MEDICATION - STANDARD ORDERS

Medications listed in this protocol may be used only for the purposes referenced by the associated ICEMA Treatment Protocol.

For Nerve Agent Antidote Kit (NAAK) or medications deployed with the ChemPack see Appendix I (Page 12).

Adenosine (Adenocard) - Adult (ALS)

Stable narrow-complex SVT or Wide complex tachycardia:

Adenosine, 6 mg rapid IVP followed immediately by 20 cc NS bolus, and Adenosine, 12 mg rapid IVP followed immediately by 20 cc NS bolus if patient does not convert. May repeat one (1) time.

Reference #s 7010, 7020, 14040

Albuterol (Proventil) Aerosolized Solution - Adult (LALS, ALS)

Albuterol, 2.5 mg nebulized, may repeat two (2) times.

Reference #s 4060, 7010, 7020, 14010, 14070

Albuterol (Proventil) Metered-Dose Inhaler (MDI) - Adult (LALS, ALS - Specialty Programs Only)

Albuterol MDI, four (4) puffs every 10 minutes for continued shortness of breath and wheezing.

Reference #s 4060, 4080, 7010, 7020, 14120, 14140, 14190

Albuterol (Proventil) - Pediatric (LALS, ALS)

Albuterol, 2.5 mg nebulized, may repeat two (2) times.

Reference #s 7010, 7020, 14120, 14140, 14190

Albuterol (Proventil) Metered-Dose Inhaler (MDI) - Pediatric (LALS, ALS - Specialty Programs Only)

Albuterol MDI, four (4) puffs every 10 minutes for continued shortness of breath and wheezing.

Reference #s 4060, 4080, 7010, 7020, 14120, 14140, 14190

Aspirin, chewable (LALS, ALS)

Aspirin, 325 mg PO chewed (one (1) adult non-enteric coated aspirin) or four (4) chewable 81 mg aspirin.

Reference #s 4060, 4080, 5010, 7010, 7020, 14240

Atropine (ALS) - Adult

Atropine, 0.5 mg IV/IO. May repeat every five (5) minutes up to a maximum of 3 mg or 0.04 mg/kg.

Organophosphate poisoning:

Atropine, 2 mg IV/IO, repeat at 2 mg increments every five (5) minutes if patient remains symptomatic.

Reference #s 4060, 4080, 7010, 7020, 13010, 14030, 14260

Atropine - Pediatric (ALS)*Organophosphate poisoning - Pediatrics less than 14 years of age:*

Atropine, 0.05 mg/kg IV/IO not to exceed adult dose of 2 mg, repeat at 0.1 mg/kg increments every five (5) minutes if patient remains symptomatic.

Reference #s 4060, 4080, 7010, 7020, 13010

Calcium Chloride - Adult (ALS)*Calcium Channel Blocker Poisonings (base hospital order only):*

Calcium Chloride, 1 gm (10 ml of a 10% solution) IV/IO.

Reference #s 5010, 7010, 7020, 13010

For cardiac arrest with suspected hypocalcemia, hyperkalemia, hypermagnesemia or calcium channel blocker poisoning (base hospital order only):

Calcium Chloride, 1 gm (10 ml of a 10% solution) IV/IO.

Reference #s 7010, 7020, 14050

Calcium Chloride - Pediatric (ALS)*Calcium Channel Blocker Poisonings (base hospital order only):*

Calcium Chloride, 20 mg/kg IV/IO over five (5) minutes.

Reference #s 7010, 7020, 13010

Dextrose - Adult (LALS, ALS)*Hypoglycemia - Adult with blood glucose less than 80 mg/dL:*

Dextrose 10% /250 ml (D10W 25 gm) IV/IO Bolus

Reference #s 4060, 4080, 5010, 7010, 7020, 8010, 13020, 13030, 14040, 14060

Dextrose - Pediatric (LALS, ALS)*Hypoglycemia - Neonates (0 - 4 weeks) with blood glucose less than 35 mg/dL or pediatric patients (more than 4 weeks) with glucose less than 60 mg/dL:*

Dextrose 10%/250 ml (D10W 25 gm) 0.5 gm/kg (5 ml/kg) IV/IO

Reference #s 5010, 7010, 7020, 13020, 13030, 14150, 14160, 14170

Diphenhydramine - Adult (ALS)

Diphenhydramine, 25 mg IV/IO

Diphenhydramine, 50 mg IM

Reference #s 4060, 4080, 7010, 7020, 13010, 14010

Diphenhydramine - Pediatric (ALS)

Allergic reaction:

2 years to 14 years Diphenhydramine, 1 mg/kg slow IV/IO, not to exceed adult dose of 25 mg, **or**

Diphenhydramine, 2 mg/kg IM not to exceed adult dose of 50 mg IM.

Reference #s 7010, 7020, 14140

Epinephrine (1 mg/ml) - Adult (LALS, ALS)

Severe Bronchospasm, Asthma Attack, Pending Respiratory Failure, Severe Allergic Reactions:

Epinephrine, 0.3 mg IM. May repeat after 15 minutes one (1) time if symptoms do not improve.

Reference # 14010

Epinephrine (0.1 mg/ml) - Adult (ALS)

For persistent severe anaphylactic reaction:

Epinephrine (0.1 mg/ml), 0.1 mg slow IVP/IO. May repeat every five (5) minutes as needed to total dosage of 0.5 mg.

Reference # 14010

Cardiac Arrest, Asystole, PEA:

Epinephrine (0.1 mg/ml), 1 mg IV/IO.

Reference #s 4060, 4080, 5010, 7010, 7020, 14010, 14050, 14260

Epinephrine (0.01 mg/ml) - Adult (ALS)

Post resuscitation, persistent profound nontraumatic shock and hypotension (Push Dose Epinephrine):

Prepare Epinephrine 0.01 mg/ml solution by mixing 9 ml of normal saline with 1 ml of Epinephrine 0.1 mg/ml in a 10 ml syringe. Administer 1 ml every one (1) to five (5) minutes titrated to maintain SBP more than 90 mm Hg.

Reference #s 4060, 4080, 5010, 7010, 7020, 11010, 14050, 14230

Epinephrine (1 mg/ml) - Pediatric (LALS, ALS)

Severe Bronchospasm, Asthma Attack, Pending Respiratory Failure, Severe Allergic Reactions:

Epinephrine, 0.01 mg/kg IM not to exceed adult dosage of 0.3 mg.

Reference #s 4060, 5010, 7010, 7020, 14120, 14140

Epinephrine (0.1 mg/ml) - Pediatric (ALS)

Anaphylactic reaction (no palpable radial pulse and depressed level of consciousness):

Epinephrine (0.1 mg/ml), 0.01 mg/kg IV/IO, no more than 0.1 mg per dose. May repeat to a maximum of 0.5 mg.

Cardiac Arrest:

1 day to 8 years Epinephrine (0.1 mg/ml), 0.01 mg/kg IV/IO (do not exceed adult dosage)

9 to 14 years Epinephrine (0.1 mg/ml), 1.0 mg IV/IO

Newborn Care:

Epinephrine (0.1 mg/ml), 0.01 mg/kg IV/IO if heart rate is less than 60 after one (1) minute after evaluating airway for hypoxia and assessing body temperature for hypothermia.

Epinephrine (0.1 mg/ml), 0.005 mg/kg IV/IO every 10 minutes for persistent hypotension as a base hospital order or in radio communication failure.

Reference # 14200

Epinephrine (0.01 mg/ml) - Pediatric (ALS)

Post resuscitation, profound shock and hypotension (Push Dose Epinephrine):

Prepare Epinephrine 0.01 mg/ml solution by mixing 9 ml of normal saline with 1 ml of Epinephrine 0.1 mg/ml in a 10 ml syringe. Administer 0.1 ml/kg (do not exceed adult dosage), every one (1) to five (5) minutes. Titrate to maintain a SBP more than 70 mm Hg.

Reference #s 5010, 7010, 7020, 11010, 14150, 14230

Fentanyl - Adult (ALS)

Chest Pain (Presumed Ischemic Origin):

Fentanyl, 50 mcg slow IV/IO over one (1) minute. May repeat every five (5) minutes titrated to pain, not to exceed 200 mcg.

Fentanyl, 100 mcg IM/IN. May repeat 50 mcg every 10 minutes titrated to pain, not to exceed 200 mcg.

Acute traumatic injuries, acute abdominal/flank pain, burn injuries, Cancer pain, Sickle Cell Crisis:

Fentanyl, 50 mcg slow IV/IO push over one (1) minute. May repeat every five (5) minutes titrated to pain, not to exceed 200 mcg IV/IO, **or**

Fentanyl, 100 mcg IM/IN. May repeat 50 mcg every 10 minutes titrated to pain, not to exceed 200 mcg.

Pacing, synchronized cardioversion:

Fentanyl, 50 mcg slow IV/IO over one (1) minute. May repeat in five (5) minutes titrated to pain, not to exceed 200 mcg.

Fentanyl, 100 mcg IN. May repeat 50 mcg every 10 minutes titrated to pain, not to exceed 200 mcg.

Reference #s 3050, 4060, 4080, 5010, 7010, 7020, 11020, 13030, 14070, 14090, 14100, 14240

Fentanyl - Pediatric (ALS)

Fentanyl, 0.5 mcg/kg slow IV/IO over one (1) minute. May repeat in five (5) minutes titrated to pain, not to exceed 100 mcg.

Fentanyl, 1 mcg/kg IM/IN, may repeat every 10 minutes titrated to pain not to exceed 200 mcg.

Reference #s 3050, 4080, 5010, 7010, 7020, 13030, 14180, 14190, 14240

Glucose - Oral - Adult (BLS, LALS, ALS)

Adult with blood glucose less than 80 mg/dL:

Glucose - Oral, one (1) tube for patients with an intact gag reflex and hypoglycemia.

Reference #s 7010, 7020, 13020, 14060, 14080, 14230

Glucose - Oral - Pediatric (BLS, LALS, ALS)

Hypoglycemia - Neonates (0 - 4 weeks) with blood glucose less than 35 mg/dL or pediatric patients (more than 4 weeks) with glucose less than 60 mg/dL:

Glucose - Oral, one (1) tube for patients with an intact gag reflex and hypoglycemia.

Reference #s 7010, 7020, 14170, 14160

Glucagon - Adult (LALS, ALS)

Glucagon, 1 mg IM/SC/IN, if unable to establish IV. May administer one (1) time only.

Beta blocker Poisoning (base hospital order only):

Glucagon, 1 mg IV/IO

Reference #s 4060, 4080, 7010, 7020, 13010, 13030, 14060

Glucagon - Pediatric (LALS, ALS)

Hypoglycemia, if unable to establish IV:

Glucagon, 0.03 mg/kg IM/IN, if unable to start an IV. May be repeated one (1) time after 20 minutes for a combined maximum dose of 1 mg.

Reference #s 7010, 7020, 13030, 14160, 14170

Beta blocker poisoning (base hospital order only):

Glucagon, 0.03 mg/kg IV/IO

Reference #'s 4060, 4080, 7010, 7020, 13010

Ipratropium Bromide (Atrovent) Inhalation Solution use with Albuterol Adult (ALS)

Atrovent, 0.5 mg nebulized. Administer one (1) dose only.

Reference #s 7010, 7020, 14010, 14070

Lidocaine - Pediatric (ALS)*Cardiac Arrest:*

1 day to 8 years Lidocaine, 1.0 mg/kg IV/IO

9 to 14 years Lidocaine, 1.0 mg/kg IV/IO

May repeat Lidocaine at 0.5 mg/kg after five (5) minutes; maximum total dose of 3 mg/kg.

*Reference #s 5010, 7010, 7020, 14150***Lidocaine 2% (Intravenous Solution) - Pediatric and Adult (ALS)***Pain associated with IO infusion:*

Lidocaine, 0.5 mg/kg slow IO push over two (2) minutes, not to exceed 40 mg total.

*Reference #s 5010, 7010, 7020, 11020***Magnesium Sulfate (ALS)***Polymorphic Ventricular Tachycardia:*

Magnesium Sulfate, 2 gm IV/IO bolus over five (5) minutes for polymorphic VT if prolonged QT is observed during sinus rhythm post-cardioversion.

Eclampsia (Seizure/Tonic/Clonic Activity):

Magnesium Sulfate, 4 gm IV/IO slow IV push over three (3) to four (4) minutes.

Magnesium Sulfate, 10 mg/min IV/IO drip to prevent continued seizures.

*Reference #s 5010, 7010, 7020, 8010, 14210**Severe Asthma/Respiratory Distress (ALS) (base hospital order only):*

Magnesium Sulfate, 2 gm slow IV drip over 20 minutes. Do not repeat.

*Reference# 14010***Magnesium Sulfate - Pediatric (ALS)***Severe Asthma/Respiratory Distress (base hospital order only):*

Magnesium Sulfate, 50 mg/kg slow IV drip over 20 minutes. Do not exceed the adult dosage of 2 gm total. Do not repeat.

*Reference # 14120***Midazolam (Versed) - Adult (ALS)***Behavioral Emergencies, with suspected excited delirium:*Midazolam, 2.5 mg IV/IO/IN. May repeat in five (5) minutes, **or**

Midazolam, 5 mg IM. May repeat in 10 minutes.

Maximum of three (3) doses using any combination of IV/IO/IM/IN may be administered. Contact base hospital for additional orders and to discuss further treatment options.

Reference # 14110

Seizure:

Midazolam, 2.5 mg IV/IO/IN. May repeat in five (5) minutes for continued seizure activity, **or**

Midazolam, 5 mg IM. May repeat in 10 minutes for continued seizure activity.

Assess patient for medication related reduced respiratory rate or hypotension.

Maximum of three (3) doses using any combination of IV/IO/IM/IN may be administered for continued seizure activity. Contact base hospital for additional orders and to discuss further treatment options.

Pacing, synchronized cardioversion:

Midazolam, 2 mg slow IV/IO push or IN

CPAP:

Midazolam, 1 mg slow IV/IO push may be administered one (1) time for anxiety related to application of CPAP. Contact base hospital for additional orders.

Reference #s 4060, 4080, 7010, 7020, 11020, 13020, 14060, 14210

Midazolam (Versed) - Pediatric (ALS)**Seizures:**

Midazolam, 0.1 mg/kg IV/IO with maximum dose 2.5 mg. May repeat Midazolam in five (5) minutes, **or**

Midazolam, 0.2 mg/kg IM/IN with maximum dose of 5 mg. May repeat Midazolam in 10 minutes for continued seizure.

Assess patient for medication related reduced respiratory rate or hypotension.

Maximum of three (3) doses using any combination of IV/IO/IM/IN may be administered for continued seizure activity. Contact base hospital for additional orders and to discuss further treatment options.

Reference #s 7010, 7020, 14170

Naloxone (Narcan) - Adult (BLS)**For resolution of respiratory depression related to suspected opiate overdose:**

Naloxone, 0.5 mg IM/IN, may repeat Naloxone 0.5 mg IM/IN every two (2) to three (3) minutes if needed.

For suspected Fentanyl overdose with respiratory depression:

Consider a loading dose of 4 mg IN Naloxone. If no signs of respiratory improvement, consider Naloxone 0.5 mg IM/IN every two (2) to three (3) minutes if needed.

Do not exceed 10 mg of Naloxone total regardless of route administered.

Reference #s 7010, 7020, 8030, 14060

Naloxone (Narcan) - Adult (LALS, ALS)

For resolution of respiratory depression related to suspected opiate overdose:

Naloxone, 0.5 mg IV/IO/IM/IN, may repeat Naloxone 0.5 mg IV/IO/IM/IN every two (2) to three (3) minutes if needed.

For suspected Fentanyl overdose with respiratory depression:

Consider a loading dose of 4 mg IN Naloxone. If no signs of respiratory improvement, consider Naloxone 0.5 mg IV/IO/IM/IN every two (2) to three (3) minutes if needed.

Do not exceed 10 mg of Naloxone total regardless of route administered.

Reference #s 4080, 7010, 7020, 14060

Naloxone (Narcan) - Pediatric (BLS)

For resolution of respiratory depression related to suspected opiate overdose:

1 day to 8 years Naloxone, 0.1 mg/kg IM/IN (do not exceed the adult dose of 0.5 mg per administration)
9 to 14 years Naloxone, 0.5 mg IM/IN

May repeat every two (2) to three (3) minutes if needed. Do not exceed the adult dosage of 10 mg total IM/IN.

Reference #s 7010, 7020, 8030, 14150, 14160

Naloxone (Narcan) - Pediatric (LALS, ALS)

For resolution of respiratory depression related to suspected opiate overdose:

1 day to 8 years Naloxone, 0.1 mg/kg IV/IO/IM/IN (do not exceed the adult dose of 0.5 mg per administration)
9 to 14 years Naloxone, 0.5 mg IV/IO/IM/IN

May repeat every two (2) to three (3) minutes if needed. Do not exceed the adult dosage of 10 mg total IV/IO/IM/IN.

Reference #s 7010, 7020, 14150, 14160

Nitroglycerin (NTG) (LALS, ALS)

Nitroglycerin, 0.4 mg sublingual/transmucosal.

One (1) every three (3) minutes as needed. May be repeated as long as patient continues to have signs of adequate tissue perfusion. **If a Right Ventricular Infarction is suspected, the use of nitrates requires base hospital contact.**

Nitroglycerin Paste, 1 inch (1 gm) transdermal, may not repeat.

Nitroglycerin sublingual is the preferred route of administration for ACS. Nitro Paste is a one (1) time dose and intended for when sublingual cannot be easily administered (i.e., CPAP).

Nitroglycerin is contraindicated if there are signs of inadequate tissue perfusion or if sexual enhancement medications have been utilized within the past 48 hours.

Reference #s 4060, 4080, 7010, 7020, 14010, 14240

Ondansetron (Zofran) - Patients four (4) years old to Adult (ALS)*Nausea/Vomiting:*

Ondansetron, 4 mg slow IV/IO/ODT

All patients four (4) to eight (8) years old: May administer a total of 4 mgs of Ondansetron prior to base hospital contact.

All patients nine (9) and older: May administer Ondansetron 4 mg; may repeat two (2) times, at 10 minute intervals, for a total of 12 mgs prior to base hospital contact.

May be used as prophylactic treatment of nausea and vomiting associated with narcotic administration.

Reference #s 4080, 7010, 7020, 14090, 14180, 14220

Oxygen (non-intubated patient per appropriate delivery device)*General Administration (Hypoxia):*

Titrate Oxygen at lowest rate required to maintain SPO₂ at 94%. Do not administer supplemental oxygen for SPO₂ more than 95%.

Chronic Obstructive Pulmonary Disease (COPD):

Titrate Oxygen at lowest rate required to maintain SPO₂ at 90%. Do not administer supplemental oxygen for SPO₂ more than 91%.

Reference #s 12010, 13010, 13020, 13030, 13050, 14010, 14020, 14030, 14040, 14060, 14070, 14090, 14120, 14130, 14140, 14160, 14170, 14180, 14190, 14200, 14210, 14220, 14230, 14240

Sodium Bicarbonate - Adult (ALS)*Tricyclic Poisoning (base hospital order only):*

Sodium Bicarbonate, 1 mEq/kg IV/IO

Reference #s 5010, 7010, 7020, 13010

For cardiac arrest with suspected metabolic acidosis, hyperkalemia or tricyclic poisoning (base hospital order only):

Sodium Bicarbonate, 50 mEq IV/IO

Reference #'s 7010, 7020, 14050

Sodium Bicarbonate - Pediatric (ALS)*Tricyclic Poisoning (base hospital order only):*

Sodium Bicarbonate, 1 mEq/kg IV/IO

Reference #'s 7010, 7020, 13010

Tranexamic Acid (TXA) - Patients 15 years of age and older (ALS)

Signs of hemorrhagic shock meeting inclusion criteria:

Administer TXA 1 gm in 50 - 100 ml of NS via IV/IO over 10 minutes. Do not administer IVP as this will cause hypotension.

Reference #s 7010, 7020, 14090

APPENDIX I**Medications for self-administration or with deployment of the ChemPack.**

Medications listed below may be used only for the purposes referenced by the associated ICEMA Treatment Protocol. Any other use, route or dose other than those listed, must be ordered in consultation with the Base Hospital physician.

Atropine - Pediatric (BLS, AEMT-Auto-injector only with training, ALS)

Known nerve agent/organophosphate poisoning with deployment of the ChemPack using:

Two (2) or more mild symptoms: Administer the weight-based dose listed below as soon as an exposure is known or strongly suspected. If severe symptoms develop after the first dose, two (2) additional doses should be repeated in rapid succession 10 minutes after the first dose; do not administer more than three (3) doses. If profound anticholinergic effects occur in the absence of excessive bronchial secretions, further doses of atropine should be withheld.

One (1) or more severe symptoms: Immediately administer (3) three weight-based doses listed below in rapid succession.

Weight-based dosing:

Less than 6.8 kg (less than 15 lbs):	0.25 mg, IM using multi-dose vial
6.8 to 18 kg (15 to 40 lbs):	0.5 mg, IM using AtroPen auto-injector
18 to 41 kg (40 to 90 lbs):	1 mg, IM using AtroPen auto-injector
More than 41 kg (more than 90 lbs):	2 mg, IM using multi-dose vial

Symptoms of insecticide or nerve agent poisoning, as provided by manufacturer in the AtroPen product labeling, to guide therapy:

Mild symptoms: Blurred vision, bradycardia, breathing difficulties, chest tightness, coughing, drooling, miosis, muscular twitching, nausea, runny nose, salivation increased, stomach cramps, tachycardia, teary eyes, tremor, vomiting, or wheezing.

Severe symptoms: Breathing difficulties (severe), confused/strange behavior, defecation (involuntary), muscular twitching/generalized weakness (severe), respiratory secretions (severe), seizure, unconsciousness, urination (involuntary).

NOTE: Infants may become drowsy or unconscious with muscle floppiness as opposed to muscle twitching.

Reference #s 11010, 13010, 13040

Diazepam (Valium) - Adult (ALS)

For seizures associated with nerve agent/organophosphate exposure ONLY with the deployment of the ChemPack:

Diazepam 10 mg (5 mg/ml) auto-injector IM (if IV is unavailable), **or**
Diazepam 2.5 mg IV

Reference # 13040

Diazepam (Valium) - Pediatric (ALS)

For seizures associated with nerve agent/organophosphate exposure ONLY with the deployment of the ChemPack:

Diazepam 0.05 mg/kg IV

Reference # 13040

Nerve Agent Antidote Kit (NAAK)/Mark I or DuoDote (containing Atropine/Pralidoxime Chloride for self-administration or with deployment of the ChemPack) - Adult

Nerve agent exposure with associated symptoms:

One (1) NAAK auto-injector IM into outer thigh. May repeat up to two (2) times every 10 to 15 minutes if symptoms persist.

Reference #s 7010, 7020, 13010, 13040



INLAND COUNTIES EMERGENCY MEDICAL AGENCY POLICY AND PROTOCOL MANUAL

Reference No. **14110R1**
Effective Date: ~~03/01/2010/01/21~~
Supersedes: ~~11/01/1803/01/20~~
Page 1 of 2

PSYCHIATRIC/BEHAVIORAL EMERGENCIES - ADULT

I. PURPOSE

To provide timely and appropriate treatment for patients that are violent, potentially violent, or who may harm themselves or others, including the potential use of restraints in the field or during transport.

II. FIELD ASSESSMENT/TREATMENT INDICATORS

Symptoms of ~~suspected excited delirium may include the physically combative, extreme dangerous~~ agitation, confusion and hallucinations, erratic behavior, profuse diaphoresis, elevated vital signs, hyperthermia, unexplained strength and endurance, and behaviors that include clothing shedding, shouting, and extreme thrashing when restrained.

This policy is not intended to negate the need for law enforcement personnel to use appropriate restraint equipment to establish scene-management control. Restraints should be applied by law enforcement whenever possible. If applied, an officer is required to remain available at the scene or during transport to remove or adjust the restraints for patient safety, per ICEMA Reference #6060 - Patient Restraints.

III. BLS INTERVENTIONS

- Approach patient in a calm and cautious manner.
- Ensure patent airway, obtain oxygen saturation and apply oxygen as needed.
- Restraint equipment must be either padded leather restraints or soft restraints (e.g., posey, velcro or seat-belt type).
- Apply four (4) point restraints as clinically indicated. Transport of a restrained patient should be in low to high Fowlers position. Never transport a patient in a prone position while restrained. Transport of a patient supine, while restrained, can affect respiratory function and constant monitoring of respiratory status is required.
- Perform cooling measures as clinically indicated.
- If suspected hypoglycemia, obtain a blood glucose.

IV. LIMITED ALS (LALS) INTERVENTIONS

- Perform activities identified in the BLS Interventions.

V. ALS INTERVENTIONS

- Perform activities identified in the BLS and LALS Interventions.
- If patient meets criteria for ~~potentially fatal and dangerous agitation-suspected excited delirium~~, administer Midazolam per ICEMA Reference #11010 - Medication - Standard Orders. Do not delay administration of Midazolam due to lack of vascular access as IM or IN is preferred in this circumstance. May repeat one (1) time using same method as first administered.

- Place on cardiac monitor. Continuous monitoring of a patient after administration of Midazolam is required.
- Obtain capnography, monitor waveform and numerical value. Apnea can be the result of the use of Midazolam and other medications.
- Once conditions are safe, establish IV.
- Base hospital may order:
 - For ~~potentially fatal and dangerous agitation suspected excited delirium~~ and suspected metabolic acidosis/hyperkalemia administer Sodium Bicarbonate, per ICEMA Reference #11010 - Medication - Standard Orders.

VI. REFERENCES

<u>Number</u>	<u>Name</u>
6060	Patient Restraints
11010	Medication - Standard Orders